



Predicted nitrogen tensions during repeated breath-hold diving in humans

Fahlman, A.¹, and Bostrom, B.²

¹Global Diving Research, Maple Ridge, BC V4R 2W7, Canada

²Department of Zoology, The University of British Columbia, 6270 University Blvd., Vancouver, BC, V6T 1Z4, Canada

Introduction

Despite increasing interest in breath-hold diving, limited research has been conducted to investigate the potential risk of decompression sickness (DCS). While repetitive breath-hold dives to 15–20m have been reported to cause DCS in humans (Paulev, 1965, 1967), no research has attempted to improve safety or to predict safe limits for this type of activity. We used a previously published mathematical model to predict N₂ tensions (PN₂) during hypothetical breath-hold dives in a human. Our objective is to investigate possible PN₂ levels during breath-hold diving and to provide simple guidelines to improve safety during breath-hold diving.

Objective

1. Determine approximate levels of N₂ in human tissues and blood during repeated shallow and single deep dives.
2. From the results, propose simple guidelines that could reduce the risk of decompression sickness.

Acknowledgement: This study was supported by private funds and was performed outside normal working hours. The dive profiles used in this study were suggested by Mr. Martin Stepanek. We would also like to thank Diana Temple for always being ready to help and Liz Montcalm-Smith for helping us print this poster.

Methods

- > We used a previously published model that estimates N₂ tension (PN₂, ATA) in mixed venous blood and 4 tissues (central circulation [heart, liver and kidney], muscle, brain, fat) during breath-hold diving (Fahlman et al., 2006).
- > We assumed a 70 kg man with a cardiac output of 6.1 · min⁻¹ and where central circulation, muscle, brain, fat and blood each were 3%, 74%, 2% and 15% and 6% of the body mass.
- > 2 dive profiles were used;
 - > 1) 30 repeated dives to 4 ATA (30 m, 98 ft) with a dive duration (DD) of 150 sec and for three different surface intervals durations (SI, 90, 150 or 300 sec).
 - > 2) A single dive to 9 ATA (80 m, 262 ft) with a DD of 210 sec.
- > We assumed that while submerged, the diver experienced a 50% reduction in cardiac output.
- > The proportion of the cardiac output to each tissue was 35% to the central circulation, 35% to muscle, 20% to brain and 10% to fat while at the surface. While diving, the same values were 35%, 30%, 30% and 5%.

Results

- Maximum venous PN₂ during 30 repeated dives to 30 m was 1.44 ATA for a SI of 300 sec and 1.69 ATA for a SI of 90 sec (Fig. 1).
- For repeated dives to 30 m, venous PN₂ during the first and sixth dive reached, respectively, 88% and 97% of the maximum estimated venous PN₂ during the entire series (Data not shown).
- Maximum venous PN₂ during a single deep dive was 3.0 ATA, or 310% higher than the surface equilibrium value (0.74 ATA, Fig. 2).




Figure 1. Mixed venous PN₂ following the last dive in a repeated dive series to 30 m (98 ft).

Figure 2. Mixed venous PN₂ following a single dive to 80 m (262 ft).

Conclusions

- Our simulations suggest that single deep or repeated shallow breath-hold dives can result in high mixed venous N₂ tensions that could possibly lead to DCS.
- A simple way to reduce excessive PN₂ and the risk of DCS symptoms is to have an SI exceeding DI by at least 2 times.

References

- Fahlman, A., Olszowska, A., Bostrom, B., Jones, D.R. 2006. Deep diving mammals: Dive behavior and circulatory adjustments contribute to bends avoidance. *Resp Physiol Neurobiol*. In press.
- Paulev, P., 1965. Decompression sickness following repeated breath-hold dives. *J. Appl. Physiol.* 20, 1028–1031.
- Paulev, P.E., 1967. Nitrogen tissue tensions following repeated breath-hold dives. *J. Appl. Physiol.* 22, 714–718.

Point of contact

Andreas Fahlman, e-mail: fahlman@zoology.ubc.ca