

Dispersal



Dispersal

What is dispersal?

Dispersal: the movement of an individual from its place of birth to the place where it reproduces or would have reproduced had it survived and found a mate (after Howard 1960)

1. Dispersal does *not* necessarily result in gene flow
2. Dispersal is *not* the same as migration
3. Dispersal and migration may interact (e.g., long distance migration could influence dispersal)
4. Dispersal is *not* the same as dispersion, which refers to the spatial distribution of individuals in a population
5. Dispersal can be discussed in the context of *ecological process* (contemporary) or *historical biogeographic events*. Historical dispersal events are difficult to study (but we also know they are important)


Migration: the spatially and temporally predictable movement of individuals between breeding and foraging habitats (Hendry *et al.* 2004)

Dispersal


At the center of arguments to explain *disjunct distributions*: dispersal versus vicariance

Vicariance: establishment of a barrier separating populations that were already present

Vicariance



Dispersal



Ancestral
population

Geographic
Isolation


Speciation

Dispersal


At the center of arguments to explain *disjunct distributions*: dispersal versus vicariance

With vicariance, both “new” regions were previously occupied by the taxon before the existence of a barrier (speciation happens after the appearance of a barrier)

Vicariance



Dispersal




Dispersal


At the center of arguments to explain *disjunct distributions*: dispersal versus vicariance

With dispersal, individuals actively colonize an area beyond a pre-existing geographic barrier to eventually form a new species

Vicariance



Dispersal



Ancestral
population

Geographic
Isolation


Speciation

Dispersal


We'll revisit this later, but you should keep in mind for your paper assignment:

What evidence or information do we need to establish whether disjunct populations were caused by a dispersal event or by a vicariance event?

Vicariance



Dispersal



Ancestral
population


Geographic
Isolation

Speciation

Dispersal

We'll revisit this later, but you should keep in mind for your paper assignment:

What evidence or information do we need to establish whether disjunct populations were caused by a dispersal event or by a vicariance event?



Dispersal

Features of dispersal as an ecological process:

- Can take place before or after first breeding
- May depend on environmental events or it may be “innate”
- May be density-dependent (or related to carrying capacity)
- Can be “condition dependent”
- Many taxa have sex-biased dispersal (e.g., birds and mammals)
- Successful reproduction by dispersers results in gene flow

Dispersal

Outline of dispersal topics:

- 1) Mechanisms of Dispersal
- 2) Barriers and Corridors
- 3) Range Expansion
- 4) Evolution of Dispersal and Philopatry

Mechanisms of Dispersal

What is a propagule?

Propagule: any part of an organism, life stage of an organism, or individual in a group of organisms that can reproduce and establish a new population

Challenges to establishing populations:

1. Sexual reproduction requires two individuals to found a population
2. Establishment of one species can be dependent on other species (i.e., trophic structure, mutualists, predator/prey, host/parasite interactions)



Mechanisms of Dispersal

Active dispersal


Vagility: the ability to disperse actively



Mechanisms of Dispersal

Passive dispersal

Pagility: the ability of an organism to disperse passively, where dispersal is dependent on some force external to the organism




Mechanisms of Dispersal

Sweepstakes dispersal

Sweepstakes dispersal: the partly stochastic dispersal of some individuals, and the establishment of remote or disjunct biota.

Over-water dispersal due to hurricanes.
On October 4, 1995, ~ 15 individuals of the green iguana appeared on eastern beaches of Anguilla in the Caribbean.

This species did not occur on the island previously, but arrived on a mat of logs and uprooted trees with large root masses.



Mechanisms of Dispersal

Sweepstakes dispersal

Sweepstakes dispersal: the partly stochastic dispersal of some individuals, and the establishment of remote or disjunct biota.

The Hawaiian hoary bat is most likely derived from migrating North American hoary bats that went astray.

The Farallon Islands (near the coastline of California) are a stopover site for NA hoary bats and would be the closest landfall (~3665 km from Hawaii)

Hawaiian hoary bat
(O-pay-ah-pay-ah)
Lasiurus cinereus semotus




Hoary bat
Lasiurus cinereus

Barriers and Corridors

Physiological Barriers

Atlantic Puffin



Distribution of species in the family Alcidae


Many species appear to be restricted to particular regions due to temperature tolerance. The avian family Alcidae (auk, puffins, murres) is restricted to cooler regions of the northern hemisphere, whereas the family Spheniscidae (penguins) is restricted to the southern hemisphere

Barriers and Corridors

Physiological Barriers

Mountain passes are greater barriers to dispersal in the tropics than in temperate regions because there is less overlap in thermal regimes experienced at low and high altitudes in the tropics.

Cold    Warm




Annual temperature variation


Barriers and Corridors

Physiological Barriers

Mountain passes are greater barriers to dispersal in the tropics than in temperate regions because there is less overlap in thermal regimes experienced at low and high altitudes in the tropics.







Barriers and Corridors



Data from diverse tropical ectotherms (e.g., fish, insects, reptiles, amphibians) suggest that tropical species living in stable aseasonal climates have:

1) narrower thermal tolerances than higher-latitude species

2) live in climates closer to their physiological optima

-  Current mean temperature
-  Current temperature range
-  Predicted mean temperature in 2100
-  Predicted temperature range in 2100

Barriers and Corridors

Corridors


Corridor: a non-selective dispersal route that permits the movement of individuals from one region to another.

Corridors should provide environments similar to that of the two source areas that it links

Corridors allow a taxonomically balanced assemblage of plants and animals to cross from one large source area to another

Barriers and Corridors

Historical Corridors




<http://jan.ucc.nau.edu/~rcb7/globaltext2.html>

280 mya (Early Permian)

Barriers and Corridors

Historical Corridors





<http://jan.ucc.nau.edu/~rcb7/globaltext2.html>

65 mya (Cretaceous-Paleogene)

Barriers and Corridors

Ecological corridors (i.e., habitat corridors)

Eight experimental deforestation landscapes at the Savannah River Site (SRS)




One of 8 landscapes of regenerating longleaf pine savanna

Examined connectivity while controlling for edge and area

Barriers and Corridors

Ecological corridors

Eight experimental deforestation landscapes at the Savannah River Site (SRS)



One of 8 landscapes of regenerating longleaf pine savanna

Examined connectivity while controlling for edge and area

Barriers and Corridors

Ecological corridors


Species richness for plants (3 groups)

Within-patch colours show increases in species richness during annual surveys (2001-2007)

Coloured strips outside patches (non-target areas) show increases relative to unconnected areas in 10-m increments from target habitat

Both patch type and distance from target patch edge were significant predictors of species richness across groups

Brudvig et al. 2009 PNAS



Barriers and Corridors


Filters

Filter: a selective or restrictive dispersal route that permits the movement of individuals with certain characteristics from one region to another.

A filtering dispersal route can lead to *asymmetric biotic exchange*

Example: the mammalian bias in the great American biotic exchange may have resulted from more NA animals being savanna-adapted (more savanna habitat in NA)

At the time of the exchange, the land bridge was continuous savanna




Barriers and Corridors

Filters

Filter: a selective or restrictive dispersal route that permits the movement of individuals with certain characteristics from one region to another.

Southern Origin	
Porcupines	
Glyptodonts	
Armadillos	
Giant ground sloths	
Opossums	
Northern Origin	
Rabbits	Mastodons
Field mice	Horses
Foxes	Tapirs
Bears	Peccaries
Raccoons	Camels
Weasels	Deer
Cats	



Range Expansion

Range expansion can follow one of two patterns:

1. Jump dispersal
2. Diffusive dispersal

Range Expansion

Jump Dispersal




Eruption on Krakatau island in 1883

Was heard 3110 km away in W. Australia

Pressure waves recorded 5 days after explosion

Ash propelled 80 km high

Anyone within 16 km of explosion would likely have gone deaf...




Range Expansion

Jump Dispersal


Species colonization for sea-, wind-, and animal-dispersed plants on Rakata Island, one of the Krakatau Islands in Indonesia.



Krakatau




- secondary forest
- casuarina woodland and grassland
- scattered casuarinas
- Saccharum* zone: very sparse clumps
- barren ash or lava
- pre-1883 island of Krakatau




Range Expansion

Jump Dispersal




Species colonization for sea-, wind-, and animal-dispersed plants on Rakata Island, one of the Krakatau Islands in Indonesia.



Range Expansion

Diffusive Dispersal





European starlings (*Sturnus vulgaris*) were introduced intentionally to North America.

The American Acclimatization Society for European settlers, which hoped to introduce to the U.S. every bird mentioned in Shakespeare's scripts, released 80-100 starlings in New York City's Central Park in 1890 and 1891.

Now a population of > 200 million

Range Expansion

Diffusive Dispersal




Northern range limit likely determined by cold tolerance and low heat conservation

The nine-banded armadillo (*Dasypus novemcinctus*) is native to South America and continues to expand its range northward.

Range Expansion

Diffusive Dispersal



Fertile Crescent crops:

- Wheat
- Barley
- Flax
- Chick peas
- Lentils

Range Expansion

Jump and Diffusive Dispersal



The cattle egret (*Bubulcus ibis*) crossed the South Atlantic under its own power and colonized northeastern South America in the late 1800s

The species then expanded its range over long periods of time via diffusive dispersal and population expansion

...many invasive species colonize and spread through a combination of jump and diffusive dispersal