

Contemplating the future: Acting now on long-term monitoring to answer 2050's questions

DAVID B. LINDENMAYER,^{1,2*} EMMA L. BURNS,^{1,2} PHILIP TENNANT,^{1,2}
CHRIS R. DICKMAN,^{1,3} PETER T. GREEN,^{1,4} DAVID A. KEITH,^{1,5}
DANIEL J. METCALFE,^{1,6} JEREMY RUSSELL-SMITH,^{1,7} GLENDA M. WARDLE,^{1,3}
DICK WILLIAMS,^{1,8} KARL BOSSARD,^{1,2} CLAIRE DELACEY,^{1,2} IVAN HANIGAN,^{1,2}
C. MICHAEL BULL,⁹ GRAEME GILLESPIE,^{1,10} RICHARD J. HOBBS,¹¹
CHARLES J. KREBS,¹² GENE E. LIKENS,^{13,14} JOHN PORTER¹⁵
AND MICHAEL VARDON²

¹Long Term Ecological Research Network, Terrestrial Ecosystem Research Network, Australia, ²Fenner School of Environment and Society, The Australian National University, Canberra, Australian Capital Territory (Email: david.lindenmayer@anu.edu.au), ³Desert Ecology Research Group, School of Biological Sciences, The University of Sydney, Sydney, New South Wales, ⁴Department of Ecology, Environment and Evolution, La Trobe University, Melbourne, Victoria, ⁵School of Biological, Earth and Environmental Sciences, The University of New South Wales, Sydney, New South Wales, ⁶CSIRO Land and Water Flagship, Dutton Park, Queensland, ⁷Darwin Centre for Bushfire Research, Charles Darwin University, Darwin, Northern Territory, ⁸Sustainable Ecosystems, CSIRO Ecosystem Sciences, Winnellie, Northern Territory, ⁹School of Biological Sciences, Flinders University, Adelaide, South Australia, ¹⁰Department of Land Resource Management, Flora and Fauna Division, Berrimah, Northern Territory, ¹¹School of Plant Biology and ARC Centre of Excellence for Environmental Decisions, University of Western Australia, Crawley, Western Australia, ¹²Institute for Applied Ecology, University of Canberra, Canberra, Australian Capital Territory, ¹³Cary Institute of Ecosystem Studies, Millbrook, New York, USA ¹⁴Department of Ecology and Environmental Biology, University of Connecticut, Storrs, Connecticut, USA and ¹⁵Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia, USA

Abstract In 2050, which aspects of ecosystem change will we regret not having measured? Long-term monitoring plays a crucial part in managing Australia's natural environment because time is a key factor underpinning changes in ecosystems. It is critical to start measuring key attributes of ecosystems – and the human and natural process affecting them – now, so that we can track the trajectory of change over time. This will facilitate informed choices about how to manage ecological changes (including interventions where they are required) and promote better understanding by 2050 of how particular ecosystems have been shaped over time. There will be considerable value in building on existing long-term monitoring programmes because this can add significantly to the temporal depth of information. The economic and social processes driving change in ecosystems are not identical in all ecosystems, so much of what is monitored (and the means by which it is monitored) will most likely target specific ecosystems or groups of ecosystems. To best understand the effects of ecosystem-specific threats and drivers, monitoring also will need to address the economic and social factors underpinning ecosystem-specific change. Therefore, robust assessments of the state of Australia's environment will be best achieved by reporting on the ecological performance of a representative sample of ecosystems over time. Political, policy and financial support to implement appropriate ecosystem-specific monitoring is a perennial problem. We suggest that the value of ecological monitoring will be demonstrable, when plot-based monitoring data make a unique and crucial contribution to Australia's ability to produce environmental accounts, environmental reports (e.g. the State of the Environment, State of the Forests) and to fulfilling reporting obligations under international agreements, such as the Convention on Biological Diversity. This paper suggests what must be done to meet Australia's ecological information needs by 2050.

Key words: ecosystem-specific monitoring, environmental accounting, environmental management, network of monitoring sites.

*Corresponding author.

Accepted for publication October 2014.

INTRODUCTION

The planet is changing rapidly as a result of increasing human population; land, ocean and climate transformations (IPCC & Editors 2013); and biodiversity loss (Butchart *et al.* 2010; Barnosky *et al.* 2012). The current rate of change is greater than at any previous period known to science (Rockström *et al.* 2009; Hooper *et al.* 2012). Many predictions are being made about the conditions likely to characterize the planet in the future (e.g. <http://hsctoolkit.bis.gov.uk/The-tools.html>; KPMG International and The Mowat Centre 2013). Some authors refer to 2050 as being a 'crunch-time' for humanity in terms of dealing with the multiple demands of a large and resource-intensive human population and rapidly dwindling natural resources (Turner 2008; Holloway 2012; Fulton 2013). But these predicted trends need to be rigorously examined so that they can be validated, adapted or dismissed (Andersen *et al.* 2014). That is, in addition to making predictions about change based on models of unknown accuracy, we also need to measure directly what is changing, how it is changing and why it is changing. This process will help track current trajectories of change relative to previous predictions and inform future predictions. It will also improve society's capacity to adapt, innovate and avoid the occurrence of predicted worst-case scenarios.

Uncertainty about the present and future state of the environment contributes substantially to the ultimate costs of addressing environmental change (Pindyck 2007; Dietz & Fankhauser 2010). It is difficult to formulate cost-effective policies to address changes that are poorly understood either with respect to magnitude of change or driving mechanisms. A relatively small amount of money spent on long-term monitoring can help better define problems and their solutions, thereby reducing the chance of expensive mishaps. Thus, recognition of change and understanding the causes of change require long-term investment in data collection. Indeed, many questions in ecology and environmental science cannot be addressed without long-term monitoring and research (Likens 1989; Muller *et al.* 2011; Lindenmayer *et al.* 2012).

In recent years in Australia, there has been growing recognition of the need to conduct environmental monitoring, with some progress made through the establishment of the *Environmental Accounting Function* within the Bureau of Meteorology under the *National Plan for Environmental Information* (Bureau of Meteorology 2014). This led to significant products such as the Biodiversity Profiling report (Zerger *et al.* 2013). This initiative, however, was curtailed in 2014, making it apparent that the basic case for environmental accounting needs to be reinvigorated. A new dialogue needs to emerge that emphasizes the importance of implementing appropriately stratified ecosystem-specific, site-

based monitoring that can detect change and explain the drivers of that change (Burns *et al.* 2014). Importantly, this approach is distinctly different from approaches to reporting on ecosystem changes, which rely heavily on a large body of inventory data (Hampton *et al.* 2013). In Australia, many of these data are now accessible through the Atlas of Living Australia or the Australian Ecological Knowledge and Observation System. Data housed within these important repositories are drawn from a variety of sources, ranging from standardized surveys undertaken by government agencies to opportunistic sightings recorded by amateur naturalists. While these repositories constitute impressive inventories in themselves, care should be taken when using these data for scientific monitoring and explaining ecological phenomena and predicting their trajectories into the future. This is because common features of such databases, such as unquantified spatial bias, the use of non-standardized sampling methods, lack of taxonomic rigour and a lack of spatial accuracy in data collection, can limit the utility of the information they contain.

Fit-for-purpose long-term ecological monitoring and research are essential if we are to answer key questions about environmental changes, particularly gradual change happening in small iterations (i.e. chronic change) rather than abrupt (acute) change resulting from a sudden alteration in conditions. However, there is a very patchy and disjunct history of long-term environmental research and monitoring in Australia (Youngentob *et al.* 2013). For example, because of a paucity of credible long-term ecological monitoring, it has been virtually impossible to tell how effective actions associated with billions of dollars of annual expenditure have been on environmental management outcomes in Australia (Hajkowicz 2009; Pannell & Roberts 2010). In addition, environmental reporting initiatives like the five-yearly State of the Environment reports (produced by the Commonwealth Department of the Environment) and the State of the Forests reports (produced by the Australian Bureau of Agricultural and Resource Economics and Sciences) are largely disconnected from any long-term ecological monitoring programmes or from other major programmes designed to improve environmental outcomes (Lindenmayer & Gibbons 2012). Instead, they are reliant on 'multiple lines of evidence', none of which is appropriately designed to provide adequate information on the condition of the environment relative to its natural fluctuations and ecosystem drivers.

In light of the problems outlined above, coupled with the suggested risks of an impending 'environmental crunch', a key overarching question is:

What should we begin measuring now that can help society better understand and manage natural

resources by 2050 (and beyond) and, in turn, guide human societies through a likely transition to a less bountiful world?

We argue that to improve natural resource management by 2050, we must (i) begin measuring key components of ecosystems systematically and purposefully now, (ii) establish the necessary infrastructure on-ground to facilitate ecological monitoring, and (iii) further develop information management architecture to archive, analyse and reuse the data at appropriate scales. This should inform the public about the status of the environment and help decision makers implement more sustainable environmental management. We outline the features that would characterize a successful nationwide monitoring initiative capable of serving the public interest towards 2050. We also summarize some of the general principles that should guide efforts to collect meaningful ecological measurements from terrestrial ecosystems. We do not make specific recommendations regarding the ecosystems and parameters to be monitored, but rather focus on general recommendations.

CHARACTERISTICS OF EFFECTIVE ECOSYSTEM MONITORING BY 2050

Prior to embarking on any credible set of ecological monitoring programmes, it is essential to properly define an ecosystem (Keith *et al.* 2013). This is to ensure that all stakeholders are working with common concepts and units for monitoring and reporting. An ecosystem is identified by four key elements: a biotic complex, an abiotic complex, the processes and interactions that link them and drive ecosystem change, and the distributional area they occupy (Keith *et al.* 2013). These elements are implicit in the System for Environmental-Economic Accounting (United Nations 2014) and also the recent IUCN process for identifying by 2025 a global Red List of Ecosystems (Keith *et al.* 2013). Such ecosystem-specific elements mean that the majority of entities to target for long-term monitoring will vary among ecosystems according to differences in ecosystem processes (including threatening processes and the interventions designed to mitigate them), differences in biota and other factors. Thus, suitable entities for long-term measurement in, for example, a desert ecosystem may well be markedly different from those in a temperate woodland. This is highlighted in a special edition of *Austral Ecology* (Nicholson *et al.* 2015), which contains a series of assessments of ecosystems in the southern hemisphere employing the IUCN Red List of Ecosystems criteria. It follows that continental reporting of the environment will be done best by detailed and focused monitoring and subsequent reporting on envi-

ronmental performance within an ensemble of targeted ecosystems over time.

In the remainder of this section, we outline the key elements that should underpin the development of robustly designed and implemented (and consequently long lasting) ecological monitoring programmes within targeted ecosystems.

1. Complete an audit of existing monitoring programmes and long-term ecological research to determine what work has been completed where and by whom (e.g. Youngentob *et al.* 2013). This is critical for taking ecological, financial and policy advantage of pre-existing long-term work with an already documented time series of information. Building greater time depth increases the potential for increased inference (Lindenmayer *et al.* 2012). This is because time can be a key variable influencing the effects of particular processes and the effectiveness of particular interventions, such as ecological restoration (Rey Benayas *et al.* 2009) and invasive species control (Buckley 2008). It is also cost-effective to build on previous research investments, depending on the research question at hand. However, there will be a need to establish new long-term monitoring to document changes within a more representative array of ecosystems, in populations of additional species or communities, or in response to additional ecological processes and management interventions (including responses to emerging environmental issues (Sutherland *et al.* 2012)). For example, some widespread and ecologically important Australian ecosystems, such as those dominated by Mitchell grass (319 000 km² across Queensland, Northern Territory, Western Australia and New South Wales (Orr & Holmes 1984)), are currently highly deficient in robust monitoring efforts, especially with respect to biodiversity responses to pastoralism (White *et al.* 2014).
2. Target environmental monitoring within a subset of key ecosystems across the Australian continent. Choosing a subset of ecosystems to robustly monitor should be guided by an appropriate stratification that leads to a range of variation in biota, physical environments and ecosystem processes being monitored nationwide. Priority ecosystems for selection should also be those suggested by standardized processes, such as evidence-based risk assessments, which could highlight those ecosystems that are most subject to threatening processes and activities, and therefore likely to benefit most from systematic experimentation and monitoring.
3. Develop standardized, evidence-based conceptual models using accepted eco-evidence

frameworks (e.g. Webb *et al.* 2011; Norris *et al.* 2012) that reflect collective understanding of ecosystem functionality (e.g. see White *et al.* 2013). Systematic synthesis of evidence will greatly improve the transparency and defensibility of decisions. A more 'evidence-based' approach to environmental management will also lead to improved environmental outcomes.

4. Identify and document the key environmental drivers in each ecosystem that require targeted monitoring. These include a range of threatening processes (which increasingly interact), such as habitat loss and fragmentation, invasive species and exotic pathogens, hunting or other kinds of harvesting, pollution, climate variability and

climate change, and human population growth (Table 1, and see Evans *et al.* 2011). We need to document and compare the relative frequency and severity of drivers of change that act as chronic pressures, such as salinity, with those that act as acute pressures, such as cyclones and severe bushfires. We also need to understand the scale at which they have impacts. In an Australian context, there are already well-developed maps and spatial prioritizations of where particular kinds of threatening processes predominate, and these can provide a valuable basis to help target monitoring (Evans *et al.* 2011). Similarly, given that climate change is likely to be a major driver and threat to ecosystems and biota per se in

Table 1. Examples of threatening processes in some Australian ecosystems based broadly on the threat classifications of Salafsky *et al.* (2008) and Auld & Keith (2009)

IUCN threat class	Threatening processes	Applicability to terrestrial Australian ecosystems
Residential and commercial development	Clearing and fragmentation	Ecosystem-specific but relevant to many non-protected areas (and some protected environments)
Agricultural and aquaculture expansion and intensification	Clearing and fragmentation	Ecosystem-specific, mainly woodlands, grasslands and wetlands
	Grazing by domestic livestock	Ecosystem-specific, typically woodlands, grasslands, shrublands and deserts
	Soil disturbance and degradation; introduction of pathogens; erosion and subsidence	Ecosystem-specific, typically woodlands, grasslands, shrublands and deserts
Energy production and mining	Exploration and mining for coal, iron ore, bauxite, gold, uranium, oil, gas	Ecosystem-specific, dependent on location of resources
Transportation and service corridors	Fragmentation	Pervasive, but more common on flat terrain
Consumptive use of 'wild' biological resources	Harvesting of species; loss of habitat	Ecosystem-specific
Human intrusions and disturbance from non-consumptive use	Bio-prospecting	Ecosystem-specific
	Tourism	Ecosystem-specific, but often relevant to protected areas
Natural system modifications (disturbance regimes)	Altered fire regimes	Pervasive in most ecosystems
	Altered hydrological regimes	Ecosystem-specific, notably wetlands and groundwater-dependent ecosystems
	Salinity	Ecosystem-specific, usually relevant to woodlands and wetlands
	Removal of dingoes	Ecosystem-specific, most evident in deserts, savanna and shrublands
Invasive and other problematic species and genes	Grazing by over-abundant native herbivores	Ecosystem-specific, usually relevant to woodlands, grasslands, shrublands and deserts
	Disease	Ecosystem-specific, for example heathlands
	Invasive predators	Pervasive in many ecosystems
	Invasive herbivores	Ecosystem-specific
	Invasive plants	Pervasive in most ecosystems
Pollution	Eutrophication	Ecosystem-specific, usually those associated with urban and agricultural areas
Geological events	Landslides	Ecosystem-specific, often on steep land
Climate change	Increased frequency and intensity of droughts, storms, heat waves, sea-level rise	Pervasive and ecosystem-specific

Australian ecosystems (Steffen *et al.* 2009), maps of where such impacts are likely to have greatest effect (Burrows *et al.* 2014) will be important for guiding where to monitor as well as what to monitor (and also how to monitor those targeted entities). A powerful way to quantify the effects of a particular ecosystem threat is to ensure that monitoring is conducted, wherever possible, not only where those threats manifest, but also where they are absent or limited.

5. Identify the important kinds of management interventions in each ecosystem that are needed or are currently implemented to mitigate the impacts of threatening processes. These interventions need to be evaluated over time to gauge the effectiveness of prescriptions, such as reservation, maintaining or enhancing ecosystem connectivity, rehabilitation, fire or grazing control, and invasive species control.
6. Select particular entities for monitoring that are likely to respond significantly to important environmental drivers, threatening processes and management interventions. These entities will be characteristic of particular ecosystems and could include ecosystem spatial extent, structural features, species composition and dominance, populations of species, and/or key ecological processes (including threatening processes). The target entities for long-term monitoring will vary among ecosystems in response to among-system differences in key ecosystem processes (including threatening processes and the interventions designed to mitigate them; see Table 1), differences in biota and other factors. This means that suitable entities for long-term monitoring in, for example, a dry sclerophyll forest ecosystem will most likely be different from those in an ephemeral wetland. Selection of target entities for monitoring should be based on several criteria, including (i) suitability for answering predefined and evolving key questions about conditions in a particular environment, (ii) the potential for (and sensitivity to) change over time, and (iii) feasibility for repeated monitoring. Feasibility for ecological monitoring does not mean a focus only on entities that are cheap to monitor, which risks directing effort away from entities crucial for answering key questions. If the target entities are elements of biodiversity, they should be a subset of biota and the abiotic component of ecological processes and interactions that influence biodiversity. This is because it is not logistically or financially possible to monitor all biodiversity. Rather than monitoring many things poorly, we should strive to monitor a few things well, as this can increase the power to reliably detect change (Lindenmayer & Likens 2010).
7. Consider additional structure in the stratified design of a long-term monitoring programme in relation to scale of ecosystem extent, with a particular focus on those parts of an ecosystem thought likely to show responses to change in important drivers (see Burrows *et al.* 2014). This approach should include stratification of sites across climatic, edaphic, latitudinal, disturbance or other gradients within an ecosystem targeted for monitoring. The use of some form of probability sampling that involves randomization to guide site selection will also provide greater confidence in generalizing results from a subset of chosen survey sites (Welsh 1996).
8. Balance monitoring effort strategically between problem-focused and surveillance-oriented approaches. Problem-focused monitoring programmes aim to improve understanding of identified environmental problems by tracking ecosystem responses under different management scenarios. When designed in a scientifically sound fashion, they are more likely than surveillance monitoring to deliver informative, cost-effective and relevant outcomes, but they may not detect responses to untargeted processes and emerging threats. Surveillance-oriented approaches may detect unexpected trends and problems. By definition, they cannot be shaped to measure particular ecological responses. Consequently, they risk poor returns on investment when no trends are detected, but may occasionally return windfalls in the form of important discoveries (e.g. long-term, pesticide-derived changes in eggshell thickness in birds (Olsen *et al.* 1993)). An appropriate balance would be a significant weighting towards problem-focused monitoring, with limited effort directed towards surveillance monitoring. Over-investment in surveillance monitoring at the expense of problem-focused monitoring is unlikely to deliver progress on the most pressing environmental imperatives (Likens & Lindenmayer 2011).
9. Recognize that continental reporting of the environment will often entail reporting on the environmental performance within particular, targeted ecosystems over time. This is because, as outlined above, ecosystem properties, characteristics, biota, drivers and threats vary markedly among ecosystems (Evans *et al.* 2011). Although a systematic approach is often required in monitoring, it is likely that those approaches will need to be varied to enable the effect of ecosystem-specific processes, functions and threats to be quantified. For example, even the same individual species may need to be monitored in different ways in different ecosystems (Sutherland 1996; Michael *et al.* 2012), have different habitat




Fig. 1. Conceptual model highlighting key linkages between monitoring and environmental reporting.

requirements in different ecosystems (Morrison *et al.* 2006) and be subject to quite different threats in those ecosystems (Lindenmayer *et al.* 2011). Therefore, many of the appropriate entities (although not all) to monitor to reflect environmental performance will vary among particular ecosystems. For example, what is sensible to monitor in the tropical savannas of northern Australia may be largely irrelevant in the temperate rainforests of south-western Tasmania (Lindenmayer *et al.* 2014). Hence, reporting (nationally and globally) under such an ecosystem-specific approach would best comprise reports on temporal trends for ecosystem extent in some ecosystems, the composition of particular communities in other ecosystems, populations of target species (such as threatened species) in others and the impacts of key ecosystem processes (e.g. altered fire regimes) in yet others, although some ecosystems might support all four broad kinds of monitoring (Lindenmayer *et al.* 2014). Accordingly, monitoring work might necessarily be conducted at different spatial scales in different ecosystems, but the common thread will be the collection of high-quality longitudinal data within a single information management system (as discussed below). This permits an interpretive synthesis of trends over time. Notably, an ecosystem-specific approach has recently been employed in a major book on Australian ecosystems, which provides a continent-wide overview of selected long-term ecological research in Australia (Lindenmayer *et al.* 2014). Where it is practicable to do so, trends identified from localized longitudinal studies may be scaled up to the ecosystem as a whole using appropriate spatio-temporal datasets. Such approaches have been undertaken in a wide array of long-term monitoring studies, including broader regional extrapolation of fire effects on biodiversity elements derived from the

- Three Parks Savanna Fire-Effects Plot Network in northern Australia (Russell-Smith *et al.* 2014).
- Entrench monitoring by linking the data streams generated from major reporting initiatives such as the State of the Environment and State of the Forests reporting; meeting international obligations under the Convention on Biological Diversity (Secretariat of the Convention on Biological Diversity 2005) emerging global policy initiatives such as the Ecosystems Red List being undertaken by the IUCN (Rodríguez *et al.* 2011; Keith *et al.* 2013); and environmental accounting (United Nations 2014) (Figs 1,2). The creation of an integrated set of environmental accounts (e.g. soil, biodiversity, carbon and water accounts) makes explicit the sources of the impacts of the economy on the environment (and vice versa). This means it becomes possible to consider the environment in regular economic planning processes (United Nations 2014; Vardon 2012). It would also enable Australia to address Aichi Target 2 of the Convention on Biological Diversity, that is: 'By 2020, at the latest, biodiversity values have been integrated into national and local development and poverty reduction strategies and planning processes and are being incorporated into national accounting, as appropriate, and reporting systems' (Secretariat of the Convention on Biological Diversity 2010).

SPECIFIC DESIGN FEATURES

There is a fundamental need to undertake and then maintain long-term ecological monitoring of targeted entities within a selection of ecosystems within a design framework that provides confidence in generalizing the results to unmonitored areas. Below, we make 10 additional general recommendations about how best to invest in, and maximize, the value of long-term ecological monitoring.




Fig. 2. The potential links between ecosystem monitoring and environmental accounting and reporting. The sequence of steps underscores the critical importance of appropriate study design and from that high-quality field-based monitoring data. These are the fundamental building blocks not only in environmental accounts, but also in predicting future conditions in an ecosystem under different management decisions and interventions. The study design, and the quality and availability of data, links directly to environmental policy and decision making.

1. Ensure that the same protocols are employed over the duration of any given long-term study. This is important to prevent confounding effects between changes in measurement methods and temporal changes in the target entities of interest. If the protocols have to be changed, then calibrate new methods of measurements with the previous methods – and document the change (including when the changes were made).
2. Establish reference plots or sites (wherever appropriate) for investigating the monitoring themes at hand. These should be adequately replicated to allow appropriate interpretation of the trends observed. The use of reference plots is essential because a key part of documenting temporal responses to threatening processes involves the quantification of responses not only in places where those processes are active but also where such processes are absent or where they have been mitigated (e.g. through management intervention) (Caughley & Gunn 1996). For example, monitoring to assess the effectiveness of reserves should be done both inside and outside protected areas (Kelaher *et al.* 2014; Rayner *et al.* 2014).
3. Measure particular targeted entities directly wherever possible, rather than measuring proxies or surrogates for that entity (see Lindenmayer & Likens 2011). For example, measure the abundance of animal species X rather than the occurrence of a particular tree species that is thought to be an indicator of animal species X. This is because the surrogacy relationship between the target entity and the proxy might not remain consistent over time or in different places (Caro 2010; Zettler *et al.* 2013).
4. Record the raw data of a given target entity, such as the presence or abundance of individual species of reptiles rather than only composite values (e.g. composite metrics like the number of species present). This is because raw values can later be aggregated to give a composite metric, but if only composite measures are gathered they cannot later be disaggregated to give raw values.
5. Understand that the frequency of temporal measurements taken is important to rigorously document trends. This is related to the variability of the system (Wilson *et al.* 2011) and is especially critical in ecosystems characterized by high temporal variability in conditions, as in many parts of Australia (McMahon *et al.* 1992). While observations taken in two periods a long time apart can be interesting, they may reveal little about trends, especially when there is considerable inter-year variability in the measured parameters (McNamara & Harding 2004; Lindenmayer & Cunningham 2011). The need for an appropriate frequency of monitoring does not specify that it must be regular or the same frequency across ecosystems. For example, more frequent measurements (within years) may be

appropriate in times of large changes compared with relatively static periods, as found in the boom and bust dynamics of desert ecosystems (Dickman *et al.* 2014).

6. Directly measure co-variables and factors that influence (or are strongly correlated with) measured response variables, such as climatic conditions or the amount of vegetation cover coincident with bird monitoring. This provides a powerful approach to document the relationships between change in a given entity (e.g. animal abundance) and the change in key attributes of the environment (e.g. the spatial extent of vegetation cover (see Cunningham *et al.* 2014)).
7. Specify meaningful trigger points within a given monitoring programme to activate key management responses well before major problems manifest, such as catastrophic declines in populations of a threatened species (Martin *et al.* 2009; Lindenmayer *et al.* 2013) or substantial increases in the impacts of an invasive plant or animal. Such trigger points, coupled with the implementation of additional management interventions that attempt to deal with these new and/or developing problems, might demand changes to monitoring designs, such as altering the frequency of monitoring (Lindenmayer & Likens 2009), although without breaching measurement protocols (if at all possible; see Point #1 above).
8. Track details about the history of plots, sites or other units that are the target for measurement in long-term monitoring programmes. This will provide context for how things have changed, which is important for diagnosing causes of change now or in 2050. But if asked now, then key aspects of history would include (i) the prior state of the system, such as conditions at the beginning of restoration (Egler 1954), (ii) the number, types and spatial patterns of biological legacies remaining after previous disturbances (Franklin *et al.* 2000; Banks *et al.* 2011), and (iii) patterns of site affinity for animals (Gill 1995). In addition to recording site history and initial site conditions, it also can be important to record other information, such as the amount and type of invasive plant and animal control, or the timing, cost and type of fencing to exclude domestic livestock. Documentation of the history of management intervention is often rare or patchy, even though interventions can have profound effects on the biota and/or ecosystem extent and condition. We suggest that documenting management interventions should include records of the amount of money spent so that the cost-effectiveness of interventions can be determined in relation to the biodiversity outcomes that have been derived. Some details about site characteristics and history should be informed by the study site stratification process.
9. Properly manage, archive and publish the datasets accumulated to make them discoverable to others (White *et al.* 2013). The efficient organization of datasets for analysis and synthesis is vital to any future use, but it is all too easy for these databases to become complicated and unwieldy. Unfortunately, training in database design is mostly restricted to computer scientists; ecologists often make do with suboptimal database solutions. Too often, poor design or lack of curation leads to potentially valuable datasets being lost or rendered virtually valueless (Pullin & Salafsky 2010). Management and archiving of datasets must include meta-data that document the way things have been measured. This will allow others to adopt comparable methods allowing them to build on past datasets and maximize the reusability of existing datasets. It will also assist data analysis and interpretation and facilitate re-analysis if new methods of data analysis and interpretation become available in the future. Capturing meta-data should also include contextual information on how a given long-term monitoring programme started, the rationale for its inception, initial objectives and underpinning methods like site selection. This information is particularly critical for long-term datasets in which the time span of data collection should extend beyond the career spans of the people responsible for instigating, establishing and implementing monitoring projects. Ideally, in 2050, we should have a readily accessible archive that is founded on, and extends, work that has documented what studies have previously been done, what studies are still current, what was measured, and what is still being measured and how (see Youngentob *et al.* 2013).
10. Recognize that curating critically important environmental datasets comes at a non-trivial cost (Berman & Cerf 2013). These costs must be factored into the budgeting for all major programmes, and individual projects, as well as the approvals for infrastructure and development projects (e.g. for the ongoing monitoring associated with mining) (Mudd 2014).

GENERAL DISCUSSION

Ecosystem-specific measurements

The selection of response variables for ecological monitoring, whether species-based or ecosystem

process-based, will depend on what is most appropriate for detecting and quantifying change in a given ecosystem (Keith *et al.* 2013). In some cases, there will be important synergies from simultaneously linking species, community and ecosystem process monitoring (Likens & Lindenmayer 2012), thereby enabling conclusions to be drawn not only about how processes influence biotic patterns, but also about how particular patterns (e.g. changes in the spatial coverage of vegetation cover) influence other patterns (such as the occurrence of species of birds) (Cunningham *et al.* 2014).

The assessment of important ecological processes in given ecosystems, including threatening processes, can be useful for identifying and quantifying what important priority actions need to be undertaken and where (Table 1). Such management actions would also then be assessed as part of monitoring programmes. Identification of priority actions can provide the basis for a continental strategy around what needs to be invested, and where, to achieve what outcomes. This can give politicians, policy makers and the general public a sense of how much funding is required to adequately address environmental problems across the continent – a national, bipartisan, whole-of-government strategy rather than a piecemeal one.

Entrenching long-term monitoring into environmental accounting

Long-term ecological monitoring has consistently been the last task to be funded and the first one cut in constrained budgets. The unreliable support of long-term ecological monitoring contrasts markedly with the long-term and entrenched support of the network of Bureau of Meteorology sites. The state of long-term ecological monitoring also contrasts markedly with the long-term monitoring of the Australian economy that has been achieved and maintained via the processes used to collect information through the System of National Accounts (Obst & Vardon 2014). Lessons from long-term economic monitoring can be applied to biodiversity and ecosystems (Vardon 2012). The production of economic accounts in Australia demands detailed economic monitoring that is undertaken primarily by the Australian Bureau of Statistics (Australian Bureau of Statistics 2013, 2014). Notably, such kinds of accounting revolutionized economic reporting and management in many nations around the world, and for example assisted with economic reconstruction following the Great Depression and the Second World War. Mandating environmental accounting has the potential, if done properly, to create the financial, logistical, legislative and governance frameworks that permanently

entrench robust programmes of long-term ecosystem and biodiversity monitoring, and integrate them with existing economic and social data used by governments, business and the general public. However, environmental accounts will only be as good as the data that go into making them. The approach we have outlined here for ecosystem monitoring will ensure the quality of much needed long-term ecological monitoring data. Moreover, enhanced monitoring capability has the potential to save large amounts of money through more effective environmental management. Environmental accounting will be one way of demonstrating this (Wentworth Group of Concerned Scientists 2008).

Mandating environmental accounts would create an information system that would enhance State of the Environment and State of the Forests reporting, and also allow the environment to be better considered in mainstream economic planning and decision making. It also would enable a framework to measure the effectiveness (including costs) of the use of natural resources. Environmental accounts enable the trade-offs between environment and economy to be clearly seen and would redress the current dominance of economic information in government and decision making. The long-term plots and sites that would form part of the monitoring and generate the data used to create biodiversity and other environmental accounts would then be acknowledged as critical parts of the nation's data infrastructure and be maintained alongside social and economic data infrastructure. Ultimately, the data from designed monitoring programmes and their use in environmental accounts will enable biodiversity and ecosystems to be recognized as equally important to the functioning of society as roads, power grids, the sewerage system and other built infrastructure.

A further strategy for entrenching environmental monitoring will be to coordinate study design, project implementation and data storage through an organizational entity charged with the responsibility for doing this. The Australian Bureau of Statistics and the Bureau of Meteorology are good examples of such organizations and are widely acknowledged as independent and non-partisan. Notably, there is currently a suite of initiatives within the Australian Bureau of Statistics linked with the development of a set of environmental accounts (e.g. Australian Bureau of Statistics 2014) according to international recognized frameworks (United Nations 2014).

CONCLUDING COMMENTS

Long-term monitoring is crucial to the conservation and management of the Australian environment. Yet environmental monitoring is rarely done in this

nation, in part because there has generally been very limited support for sustained long-term ecological monitoring programmes and coordination within and across programmes. Serious environmental problems associated with resource use and management are already evident and well documented. The year 2050 is forecast as a crisis point when the consequences of past practices, in concert with continued population growth, will see the breakdown of critical ecosystem services, resulting in a less predictable and bountiful environment (Turner 2008; Holloway 2012; Fulton 2013). We must begin measuring the key components of ecosystems now and continue that work for many decades to improve ecosystem integrity and ecologically sustainable resource management. We have outlined a series of key attributes that must characterize effective ecological monitoring. These include recognition that the entities being measured, and the approaches to monitor them, will be ecosystem-specific and relevant to the key ecological processes, threatening processes, and management interventions in particular ecosystems. Strategies crucial to success include the integration of ecosystem-specific monitoring approaches with initiatives like State of the Environment reporting and systems of national environmental accounting, and the development of appropriate information management architecture. A way forward would be for the community of ecological scientists and managers to agree on a set of general principles for long-term ecological monitoring, possibly including recommendations for a new body analogous to the Australian Bureau of Statistics or the Bureau of Meteorology to coordinate long-term ecological monitoring in Australia. Indeed, this is one of the key recommendations of the plan for Australian ecosystem science (Andersen *et al.* 2014).

ACKNOWLEDGEMENTS

This paper is largely possible because of the advent of the Long Term Ecological Research Network (LTERN). LTERN is a facility within the Terrestrial Ecosystem Research Network (TERN). TERN is supported by the Australian Government through the National Collaborative Research Infrastructure Strategy. We also thank the other organizations that have supported research and monitoring over prolonged periods.

REFERENCES

Andersen A., Beringer J., Bull C. M. *et al.* (2014) Foundations for the future: a long-term plan for Australian ecosystem science. *Austral Ecol.* **39**, 739–48.

- Auld T. & Keith D. A. (2009) Dealing with threats: integrating science and management. *Ecol. Manag. Restor.* **10**, S79–87.
- Australian Bureau of Statistics (2013) *Australian System of National Accounts: Concepts, Sources and Methods*. Australian Bureau of Statistics, Canberra.
- Australian Bureau of Statistics (2014) *Australian Environmental-Economic Accounts*. Australian Bureau of Statistics, Canberra.
- Banks S. C., Dujardin M. & McBurney L. (2011) Starting points for small mammal population recovery after wildfire: recolonisation or residual populations? *Oikos* **120**, 26–37.
- Barnosky A. D., Hadly E. A., Bascompte J. *et al.* (2012) Approaching a state shift in the Earth's biosphere. *Nature* **486**, 52–8.
- Berman F. & Cerf V. (2013) Who will pay for public access to research data? *Science* **341**, 616–17.
- Buckley Y. M. (2008) The role of research for integrated management of invasive species, invaded landscapes and communities. *J. Appl. Ecol.* **45**, 397–402.
- Bureau of Meteorology (2014) *National Plan for Environmental Information Initiative*. Bureau of Meteorology, Canberra. Available from URL: <http://www.bom.gov.au/environment/about.shtml>
- Burns E. L., Lindenmayer D., Tennant P. *et al.* (2014) *Making Ecological Monitoring Successful: Insights and Lessons from the Long Term Ecological Research Network*. Biotext, Canberra. Available from URL: www.tern.org.au/ltern
- Burrows M. T., Schoeman D. S., Richardson A. J. *et al.* (2014) Geographical limits to species-range shifts are suggested by climate velocity. *Nature* **507**, 492–5.
- Butchart S. H. M., Walpole M., Collen B. *et al.* (2010) Global biodiversity: indicators of recent declines. *Science* **32**, 1164–8.
- Caro T. (2010) *Conservation by Proxy. Indicator, Umbrella, Keystone, Flagship, and Other Surrogate Species*. Island Press, Washington, DC.
- Caughley G. C. & Gunn A. (1996) *Conservation Biology in Theory and Practice*. Blackwell Science, Cambridge.
- Cunningham R. B., Lindenmayer D. B., Crane M. *et al.* (2014) The law of diminishing returns: woodland birds respond to native vegetation cover at multiple spatial scales and over time. *Divers. Distrib.* **20**, 59–71.
- Dickman C., Wardle G., Foulkes J. & de Preu N. (2014) Desert complex environments. In: *Biodiversity and Environmental Change: Monitoring, Challenges and Direction* (eds D. B. Lindenmayer, E. Burns, N. Thurgate & A. Lowe) pp. 379–438. CSIRO Publishing, Melbourne.
- Dietz S. & Fankhauser S. (2010) Environmental prices, uncertainty, and learning. *Oxford Rev. Econ. Policy* **26**, 270–84.
- Egler F. E. (1954) Vegetation science concepts. I. Initial floristic composition – a factor in old field vegetation development. *Vegetatio* **4**, 412–17.
- Evans M. C., Watson J. E., Fuller R. A. *et al.* (2011) The spatial distribution of threats to species in Australia. *Bioscience* **61**, 281–9.
- Franklin J. F., Lindenmayer D. B., MacMahon J. A. *et al.* (2000) Threads of continuity. *Conserv. Practice* **1**, 8–16.
- Fulton B. (2013) Where is Australia headed? Some future projections. The Conversation. [Cited 5 November 2013.] Available from URL: <http://theconversation.com/where-is-australia-headed-some-future-projections-12403>.

- Gill F. B. (1995) *Ornithology*. W.H. Freeman and Company, New York.
- Hajkowicz S. (2009) The evolution of Australia's natural resource management programs: towards improved targeting and evaluation of investments. *Land Use Policy* **26**, 471–8.
- Hampton S. E., Strasser C. A., Tewksbury J. T. *et al.* (2013) Big data and the future of ecology. *Front. Ecol. Environ.* **1**, 156–62.
- Holloway J. (2012) Hot, crowded, and running out of fuel: earth of 2050 a scary place. [Cited 1 November 2013.] Available from URL: <http://arstechnica.com/science/2012/03/hot-crowded-and-running-out-of-fuel-earth-of-2050-a-scary-place/>
- Hooper D. U., Adair E. C., Cardinale B. J. *et al.* (2012) A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* **486**, 105–8.
- IPCC & Editors (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report in the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, USA.
- Keith D. A., Rodriguez J. P., Rodriguez-Clark K. M. *et al.* (2013) Scientific foundations for an IUCN Red List of Ecosystems. *PLoS ONE* **8**, e62111.
- Kelahr B. P., Coleman M. A., Broad A. *et al.* (2014) Changes in fish assemblages following the establishment of a network of no-take marine reserves and partially-protected areas. *PLoS ONE* **9**, e85825.
- KPMG International and The Mowat Centre (2013) *The Future State 2030*. KPMG International and The Mowat Centre, School of Public Policy and Governance, University of Toronto, Toronto.
- Likens G. E., ed. (1989) *Long-Term Studies in Ecology: Approaches and Alternatives*. Springer-Verlag, New York.
- Likens G. E. & Lindenmayer D. B. (2011) A strategic plan for an Australian Long-Term Environmental Monitoring network. *Austral Ecol.* **36**, 1–8.
- Likens G. E. & Lindenmayer D. B. (2012) Integrating approaches leads to more effective conservation of biodiversity. *Biodivers. Conserv.* **21**, 3323–41.
- Lindenmayer D. B., Burns E., Thurgate N. & Lowe A., eds (2014) *Biodiversity and Environmental Change: Monitoring, Challenges and Direction*. CSIRO Publishing, Melbourne.
- Lindenmayer D. B. & Cunningham R. B. (2011) Longitudinal patterns in bird reporting rates in a threatened ecosystem: is change regionally consistent? *Biol. Conserv.* **144**, 430–40.
- Lindenmayer D. B. & Gibbons P., eds (2012) *Biodiversity Monitoring in Australia*. CSIRO Publishing, Melbourne.
- Lindenmayer D. B. & Likens G. E. (2009) Adaptive monitoring: a new paradigm for long-term research and monitoring. *Trends Ecol. Evol.* **24**, 482–6.
- Lindenmayer D. B. & Likens G. E. (2010) *Effective Ecological Monitoring*. CSIRO Publishing and Earthscan, Melbourne and London.
- Lindenmayer D. B. & Likens G. E. (2011) Direct measurement versus surrogate indicator species for evaluating environmental change and biodiversity loss. *Ecosystems* **14**, 47–59.
- Lindenmayer D. B., Likens G. E., Andersen A. *et al.* (2012) Value of long-term ecological studies. *Austral Ecol.* **37**, 745–57.
- Lindenmayer D. B., Piggott M. & Wintle B. (2013) Counting the books while the library burns: why conservation monitoring programs need a plan for action. *Front. Ecol. Environ.* **11**, 549–55.
- Lindenmayer D. B., Wood J. T., McBurney L. *et al.* (2011) How to make a common species rare: a case against conservation complacency. *Biol. Conserv.* **144**, 1663–72.
- McMahon T. A., Finlayson B. L., Haines A. T. & Srikanthan R. (1992) *Global Runoff – Continental Comparisons of Annual Flows and Peak Discharges*. Catena Verlag, Cremlingen.
- McNamara J. M. & Harding K. C. (2004) Measurement error and estimates of population extinction risk. *Ecol. Lett.* **7**, 16–20.
- Martin J., Runge M. C., Nichols J. D., Lubow B. C. & Kendall W. L. (2009) Structured decision making as a conceptual framework to identify thresholds for conservation and management. *Ecol. Appl.* **19**, 1079–90.
- Michael D. R., Cunningham R. B., Donnelly C. F. & Lindenmayer D. B. (2012) Comparative use of active searches and artificial refuges to survey reptiles in temperate eucalypt woodlands. *Wildl. Res.* **39**, 149–62.
- Morrison M. L., Marcot B. G. & Mannan R. W. (2006) *Wildlife-Habitat Relationships. Concepts and Applications*. Island Press, Washington, DC.
- Mudd G. (2014) Mining. In: *Ten Commitments Revisited: Securing Australia's Future Environment* (eds D. B. Lindenmayer, S. Dovers & S. Morton) pp. 157–64. CSIRO Publishing, Melbourne.
- Muller F., Baessler C., Schubert H. & Klotz S., eds (2011) *Long-Term Ecological Research. Between Theory and Application*. Springer, Dordrecht.
- Nicholson E., Regan T. J., Auld T. D. *et al.* (2015) Towards consistency, rigour and compatibility of risk assessments for ecosystems and ecological communities. *Austral Ecol.*
- Norris R. H., Webb J. A., Nichols S. J., Stewardson M. J. & Harrison E. T. (2012) Analyzing cause and effect in environmental assessments: using weighted evidence from the literature. *Freshwater Sci.* **31**, 5–21.
- Obst C. & Vardon M. (2014) Recording environmental assets in the national accounts. *Oxford Rev. Econ. Policy* **30**, 126–44.
- Olsen P., Fuller P. & Marples T. G. (1993) Pesticide-related eggshell thinning in Australian raptors. *Emu* **93**, 1–11.
- Orr D. M. & Holmes W. E. (1984) Mitchell grasslands. In: *Management of Australia's Rangelands* (eds G. N. Harrington, A. D. Wilson & M. D. Young) pp. 241–54. CSIRO, Melbourne.
- Pannell D. J. & Roberts A. M. (2010) Australia's national action plan for salinity and water quality: a retrospective assessment. *Aust. J. Agric. Resour. Econ.* **54**, 437–56.
- Pindyck R. S. (2007) Uncertainty in environmental economics. *Rev. Environ. Econ. Policy* **1**, 45–65.
- Pullin A. S. & Salafsky N. (2010) Save the whales? Save the rainforest? Save the data! *Conserv. Biol.* **24**, 915–17.
- Rayner L., Lindenmayer D. B., Wood J. T., Gibbons P. & Manning A. D. (2014) Are protected areas maintaining bird diversity? *Ecography* **37**, 43–53.
- Rey Benayas J. M., Newton A. C., Diaz A. & Bullock J. M. (2009) Enhancement of biodiversity and ecosystem services by ecological restoration: a meta-analysis. *Science* **325**, 1112–24.
- Rockström J., Steffen W., Noone K. *et al.* (2009) Planetary boundaries: exploring the safe operating space for humanity.

- Ecol. Soc.* **14**, 32. Available from URL: <http://www.ecologyandsociety.org/vol14/iss32/art32/>
- Rodríguez J. P., Rodríguez-Clark K. M., Baillie J. E. M. *et al.* (2011) Establishing IUCN Red List Criteria for threatened ecosystems. *Conserv. Biol.* **25**, 21–9.
- Russell-Smith J., Edwards A., Woinarski J. C. *et al.* (2014) Northern Australia tropical savannas: the Three Parks Savanna Fire-effects Plot Network. In: *Biodiversity and Environmental Change: Monitoring, Challenges and Direction* (eds D. B. Lindenmayer, E. Burns, N. Thurgate & A. Lowe) pp. 335–78. CSIRO Publishing, Melbourne.
- Salafsky N., Salzer D., Stattersfield A. J. *et al.* (2008) A standard lexicon for biodiversity conservation: unified classifications of threats and actions. *Conserv. Biol.* **22**, 897–911.
- Secretariat of the Convention on Biological Diversity (2005) Handbook of the Convention on Biological Diversity Including Its Cartagena Protocol on Biosafety, 3rd edition. UNEP, Montreal, Canada.
- Secretariat of the Convention on Biological Diversity (2010) The Aichi Biodiversity Targets. In: *Strategic Plan for Biodiversity 2011–2020 and the Aichi Targets*. “Living Harmony with nature”. *The Strategic Plan for Biodiversity 2011–2020 – A ten-year framework for action by all countries and stakeholders to save biodiversity and enhance its benefits for people*. UNEP, Montreal, Canada. [Accessed 9 December 2014.] Available from URL: <http://www.cbd.int/doc/strategic-plan/2011-2020/Aichi-Targets-EN.pdf>
- Steffen W., Burbidge A., Hughes L. *et al.* (2009) *Australia’s Biodiversity and Climate Change*. CSIRO Publishing, Melbourne.
- Sutherland W. (1996) *Ecological Census Techniques*. Cambridge University Press, Cambridge.
- Sutherland W. J., Aveling R., Bennun L. *et al.* (2012) A horizon scan of global conservation issues for 2012. *Trends Ecol. Evol.* **27**, 12–18.
- Turner G. (2008) A Comparison of The Limits to Growth with Thirty Years of Reality. Socio-Economics and the Environment in Discussion (SEED). CSIRO Working Papers Series Number 2008–09. June 2008. ISSN 1834–5638. 52 pp.
- United Nations (2014) *System of Environmental-Economic Accounting (2012) Central Framework*. United Nations, New York.
- Vardon M. (2012) Lessons from environmental accounting for improving biodiversity monitoring. In: *Biodiversity Monitoring in Australia* (eds D. B. Lindenmayer & P. Gibbons) pp. 79–88. CSIRO Publishing, Melbourne.
- Webb J. A., Wealands S. R., Lea P. *et al.* (2011) Eco evidence: using the scientific literature to inform evidence-based decision making in environmental management. In: *MODSIM2011, 19th International Congress on Modelling and Simulation* (eds F. Chan, D. Marinova & R. S. Anderssen) Modelling and Simulation Society of Australia and New Zealand, Perth.
- Welsh A. (1996) *Aspects of Statistical Inference*. John Wiley and Sons, New York.
- Wentworth Group of Concerned Scientists (2008) *Accounting for Nature. A Model for Building the National Environmental Accounts of Australia*. Wentworth Group of Concerned Scientists, Sydney.
- White A., Orr D. M., Novelly P. & Bastin G. (2014) Tussock grassland. In: *Biodiversity and Environmental Change: Monitoring, Challenges and Direction* (eds D. B. Lindenmayer, E. Burns, N. Thurgate & A. Lowe) pp. 479–517. CSIRO Publishing, Melbourne.
- White E. P., Baldrige E., Brym Z. T. *et al.* (2013) Nine simple ways to make it easier to (re)use your data. *Ideas Ecol. Evol.* **6**, 1–10.
- Wilson H. B., Kendall B. E. & Possingham H. P. (2011) Variability in population abundance and the classification of extinction risk. *Conserv. Biol.* **25**, 747–57.
- Youngtob K., Likens G. E., Williams J. E. & Lindenmayer D. B. (2013) A survey of long-term terrestrial ecology studies in Australia. *Austral Ecol.* **38**, 365–73.
- Zerger A., Williams K. J., Nicholls M. *et al.* (2013) *Biodiversity Profiling: Components of a Continental Biodiversity Information Capability*. Environmental Information Program Publication Series No. 2, Bureau of Meteorology, Canberra, Australia.
- Zettler M. L., Proffitt C. E., Darr A. *et al.* (2013) On the myths of indicator species: issues and further consideration in the use of static concepts for ecological applications. *PLoS ONE* **8**, e78219.