

UBC Physics 102

Lecture 6

Rik Blok


<https://www.zoology.ubc.ca/~rikblok/phys102/lecture/>

UBC Physics 102 Lecture 6, July 9, 2020—p. 1/14



Electric flux [Text: Sect. 22-1]

- **Discussion: Motivation**
 - In principle can always use Coulomb's law to find field.
 - But can get very difficult. Gauss's law often easier.
- **Definition: electric flux**



<https://www.zoology.ubc.ca/~rikblok/phys102/lecture/>

UBC Physics 102 Lecture 6, July 9, 2020—p. 3/14

Outline

- ▷ Electric flux
- ▷ Gauss's law
- ▷ Applications
- ▷ End

<https://www.zoology.ubc.ca/~rikblok/phys102/lecture/>

UBC Physics 102 Lecture 6, July 9, 2020—p. 2/14



Electric flux, contd

- **Definition: electric flux, contd**
 - Flux is a measure of how much field “flows” through the surface.
 - Product of perpendicular component of electric field through a surface and the surface area,
$$\Phi_E = E_{\perp} A = E \cos \theta A.$$
 - Flux proportional to number of field lines that go through surface.
 - Magnitude of E_{\perp} must be constant over surface.
- **Interactive Quiz: PRS 06a**

<https://www.zoology.ubc.ca/~rikblok/phys102/lecture/>

UBC Physics 102 Lecture 6, July 9, 2020—p. 4/14



Electric flux, contd

- **Discussion: Closed surfaces**
 - Gauss's law deals with closed surfaces which enclose a volume of space.
 - For closed surfaces the net flux is the sum of the flux over each surface,
$$\Phi_E = \sum_{\text{surfaces}} E_{\perp} A.$$
 - Convention: E_{\perp} is +ve coming out of volume, -ve going in.

- **Interactive Quiz: PRS 06b**



Electric flux, contd

- **Discussion: Field lines**
 - If every field line entering volume also leaves then flux is always zero (regardless of shape).
 - Flux only nonzero if field lines start or stop in volume.
 - Field lines only start/stop on charges.
 - So nonzero flux means volume encloses charge.



Gauss's law [Text: Sect. 22-2]

- **Definition: Gauss's law**
 - If a closed (Gaussian) surface contains a charge Q_{encl} then

$$\sum_{\text{surfaces}} E_{\perp} A = \frac{Q_{\text{encl}}}{\epsilon_0}.$$

- Q_{encl} is sum total of all charge inside surface.
- Q_{encl} doesn't depend on positions or configuration of charges or on any charges outside surface (but electric field does).



Gauss's law, contd

- **Discussion: Gaussian surface**
 - Want to choose surface that makes problem easy to solve (usually trying to solve for E).
 - Need to know direction of E field.
 - Use symmetry.
 - Construct surface so field is parallel or perpendicular to it *everywhere*.
 - Parallel sections have zero flux, $\Phi_E = 0$.
 - Perpendicular sections have constant E_{\perp} so $\Phi_E = E_{\perp} A$ is easy to compute.



Gauss's law, contd

Derivation: Coulomb's law

- Can derive Coulomb's law from Gauss's law. Want to determine electric field at a distance r from a point charge Q .



- System looks the same from any angle so electric field must also look the same from any angle (spherical symmetry).
- E -field must point radially away from (or to) charge.
- So pick Gaussian surface so that E -field is always perpendicular: a *sphere*.




<https://www.zoology.ubc.ca/~f11k10k/phys102/lecture/>

UBC Physics 102: Lecture 6, July 9, 2008 - p. 9/14

Gauss's law, contd

Derivation: Coulomb's law, contd



- Enclosed charge: $Q_{\text{encl}} = Q$.
- Flux: $\Phi_E = E_{\perp} A = E(4\pi r^2)$ (surface area of sphere).
- Gauss's law: $Q/\epsilon_0 = E(4\pi r^2)$ or

$$E = \frac{Q}{4\pi\epsilon_0 r^2} = \frac{kQ}{r^2}. \quad \square$$



<https://www.zoology.ubc.ca/~f11k10k/phys102/lecture/>

UBC Physics 102: Lecture 6, July 9, 2008 - p. 10/14

Applications [Text: Sect. 22-3]

Discussion: Gauss's law

- Gauss's law more general than Coulomb's.
- Gauss's law one of Maxwell's equations: 4 laws that describe *all* electricity and magnetism.

Interactive Quiz: PRS 06c

Example: Long, uniformly charged line

- A very long, straight wire possesses a uniform charge per unit length λ . Calculate the electric field at a distance r from the wire.




<https://www.zoology.ubc.ca/~f11k10k/phys102/lecture/>

UBC Physics 102: Lecture 6, July 9, 2008 - p. 11/14

Applications, contd

Solution: Long, uniformly charged line

- By symmetry, the electric field must point radially out from the wire.
- So a good choice for a Gaussian surface is a cylinder (length l , radius r).



- Then $E \perp$ to side of cylinder and \parallel to ends.
- Enclosed charge: $Q_{\text{encl}} = \lambda l$.



<https://www.zoology.ubc.ca/~f11k10k/phys102/lecture/>

UBC Physics 102: Lecture 6, July 9, 2008 - p. 12/14

Applications, contd

• Solution: Long, uniformly charged line, contd

- Flux: $(E_{\perp}A)_{\text{side}} = E 2\pi r l$, $(E_{\perp}A)_{\text{ends}} = 0$.
- Gauss's law: $\sum E_{\perp}A = Q_{\text{encl}}/\epsilon_0$,

$$E 2\pi r l = \frac{\lambda l}{\epsilon_0}$$
$$E = \frac{\lambda}{2\pi\epsilon_0 r}.$$

- Could have solved with Coulomb's law but would have had to integrate.



End

• Practice Problems:

- Ch. 22: Q. 1, 2, 3, 4, 5, 7, 9, 13.
- Ch. 22: Pr. 3, 5, 7, 9, 17, 21, 23, 25, 27, 29, 31, 33, 39, 41, 43, 45, 47, 49.

• Interactive Quiz: Feedback

• Tutorial Question: tut06

