



Do extreme postprandial levels of oxygen, carbon dioxide, and ammonia in the digestive tract equilibrate with the bloodstream in the freshwater rainbow trout (*Oncorhynchus mykiss*)?

Ellen H. Jung¹ · Colin J. Brauner¹ · Chris M. Wood¹

Received: 18 February 2022 / Revised: 20 December 2022 / Accepted: 6 January 2023
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

The gastrointestinal tract (GIT) lumen of teleosts harbors extreme conditions, especially after feeding: high PCO_2 (20–115 Torr), total ammonia (415–3710 μM), PNH_3 (79–1760 μTorr in the intestine), and virtual anoxia ($\text{PO}_2 < 1$ Torr). These levels could be dangerous if they were to equilibrate with the bloodstream. Thus, we investigated the potential equilibration of O_2 , CO_2 , and ammonia across the GIT epithelia in freshwater rainbow trout by monitoring postprandial arterial and venous blood gases in vivo and in situ. In vivo blood was sampled from the indwelling catheters in the dorsal aorta (DA) and subintestinal vein (SIV) draining the posterior intestine in the fasting state and at 4 to 48 h following catheter-feeding. To investigate possible ammonia absorption in the anterior part of the GIT, blood was sampled from the DA, SIV and hepatic portal vein (HPV) from anaesthetized fish in situ following voluntary feeding. We found minimal equilibration of all three gases between the GIT lumen and the SIV blood, with the latter maintaining pre-feeding levels ($\text{PO}_2 = 25\text{--}49$ Torr, $\text{PCO}_2 = 6\text{--}8$ Torr, and total ammonia = 117–134 μM and $\text{PNH}_3 = 13\text{--}30$ μTorr at 48 h post-feeding). In contrast to the SIV, we found that the HPV total ammonia more than doubled 24 h after feeding (128 to 297 μM), indicative of absorption in the anterior GIT. Overall, the GIT epithelia of trout, although specialized for absorption, prevent dangerous levels of PO_2 , PCO_2 and ammonia from equilibrating with the blood circulation.

Keywords Digestion · Dorsal aorta · Subintestinal vein · Hepatic portal vein · Lumen · Alkaline tide

Introduction

Conditions to improve fish health and maximize growth efficiency have been of growing interest in support of aquaculture, which is currently one of the fastest expanding industries globally (Food and Agriculture Organization 2020). Despite such interest, relatively little is known about the conditions in the gastrointestinal tract (GIT) and their significance to fish productivity. The GIT is an important

multifunctioning organ in fish not only for digestion and nutrient absorption, but also for ionic and osmotic regulation, neuroendocrine regulation, immune responses, and even air breathing in some species (Grosell et al. 2011).

The GIT lumen is a unique extracorporeal environment containing a complex microbiome and lined with specialized GIT cells (e.g., oxynticopeptic cells, mucocytes, various enterocytes) interacting closely with other organ systems. In mammals, the lumen contains high carbon dioxide (CO_2) and low oxygen (O_2) levels derived from chemical reactions and bacterial fermentation of dietary substrates during digestion (reviewed by Kalantar-Zadeh et al. 2019). Recent research has shown that this is similar in fish. Direct measurements in the lumen of fasted freshwater rainbow trout (*Oncorhynchus mykiss*) and seawater English sole (*Parophrys vetulus*) have revealed low PO_2 (< 1 Torr) and high in PCO_2 (7–17 Torr) values (Wood and Eom 2019; Jung et al. 2020). Note that 1 Torr = 1 mm Hg = 0.13332 kPa. Feeding further elevates the luminal PCO_2 in fish (20–115 Torr), whereas PO_2 remains nearly anoxic (Wood and Eom 2019; Jung

Communicated by B. Pelster.

✉ Ellen H. Jung
jung@zoology.ubc.ca
Colin J. Brauner
brauner@zoology.ubc.ca
Chris M. Wood
woodcm@zoology.ubc.ca

¹ Department of Zoology, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

et al. 2020, 2022a). These luminal PO_2 and PCO_2 levels are much more extreme than normal arterial blood levels (PO_2 of ≥ 90 Torr; PCO_2 of ≤ 3 Torr, e.g. (Tetens and Lykkeboe 1985; Perry and Reid 1992; Thomas et al. 1994), and even more extreme than levels considered harmful in the natural environment (Alabaster et al. 1957; Ellis et al. 2017). Moreover, ammonia is also high in the lumen (415–3710 μM as total ammonia; 79–1760 μTorr in the intestine as PNH_3) as a result of breakdown of proteins in the feed (Bucking and Wood 2012; Rubino et al. 2014; Jung et al. 2022a). This also exceeds the water ammonia levels considered toxic for fish (Solbé and Shurben 1989; Randall and Tsui 2002).

Ultimately, there are significant gradients of both PCO_2 and ammonia from the lumen into the arterial blood perfusing the GIT, while the opposite is true for PO_2 . Notably, plasma ammonia concentrations in mixed caudal blood (Kaushik and de Oliva Teles 1985; Bucking and Wood 2008), and in the hepatic portal vein (HPV) draining the GIT (Karlsson et al. 2006) are elevated significantly after feeding. This excess ammonia is then excreted via the gills, and to a lesser extent via the kidney (Beamish and Thomas 1984; Wicks and Randall 2002; Karlsson et al. 2006; Bucking et al. 2010). However, transport of luminal ammonia into the systemic bloodstream must be carefully controlled as it can be a toxicant (Randall and Tsui 2002). Similarly, if the high luminal PCO_2 were to equilibrate with the blood, it would result in a very unfavourable blood acid–base disturbance (extreme respiratory acidosis) in fish (Perry and Gilmour 2006), though at the same time it potentially could help to release O_2 from haemoglobin by Bohr and Root effects (Nikinmaa 2006; Rummer and Brauner 2015). In mammals, some of the luminal CO_2 and other gases can be absorbed across the intestinal epithelia and enter the blood circulation, but they are also either removed by microbes or released as flatulence (Ohashi et al. 2007; Lacy et al. 2011). Whether or not high luminal PCO_2 and ammonia are also moving into the vascular system in fish is still largely unproven. A few calculated PCO_2 values in the venous blood draining from the GIT are reported to be low (Eliason et al. 2007; Cooper et al. 2014), but direct measurements in the subintestinal vein (SIV) have proven to be quite variable (Wood and Eom 2019; Jung et al. 2020, 2022a). This variability may reflect the fact that these direct measurements were taken from anaesthetized, artificially ventilated fish by direct micro-optode puncture of the SIV (in situ) rather than from conscious animals. Anaesthesia and/or blockade of blood flow may have confounded the results.

Therefore, the purpose of this study was to investigate the degree of equilibration of O_2 , CO_2 , and ammonia, between the GIT lumen and systemic bloodstream, using in vivo measurements on non-anaesthetized fish fitted with a stomach feeding tube and indwelling catheters for sampling inflowing blood from the dorsal aorta (DA) and

outflowing blood from the subintestinal vein (SIV). We investigated this in freshwater-acclimated rainbow trout sampled in the fasting state and at 4, 8, 12, 24, and 48 h after feeding (in vivo series; 1% body mass ration). We extended our in vivo ammonia findings with in situ experiments on anaesthetized trout that permitted sampling of blood from both the SIV that drains only the posterior tract and the hepatic portal vein (HPV) which additionally drains the entire GIT (in situ series; voluntarily fed to satiation). Plasma ammonia concentrations at the two sites were compared during the fasting state and at 24 and 48 h after voluntarily feeding (in situ series). In general, we hypothesized there would be minimal equilibration of these gases across the intestinal epithelia into the venous drainage. We also hypothesized that feeding would have little effect, so that SIV PO_2 , PCO_2 and ammonia would remain largely unchanged during digestion. Finally, in light of the findings of Karlsson et al. (2006), we hypothesized that postprandial luminal absorption of ammonia would be seen in the anterior part of the GIT, and thus higher plasma ammonia concentrations in blood sampled from the HPV than from the SIV.

Materials and methods

Experimental animals

Rainbow trout (220–620 g) were obtained from Little Cedar Falls Hatchery (Nanaimo, BC, Canada) and transferred to the University of British Columbia (UBC) where they were held for several months prior to experiments. Fish were held at 9°C in flowing dechlorinated Vancouver tap water ($\text{Na}^+ = 0.09$, $\text{Cl}^- = 0.10$, $\text{Ca}^{2+} = 0.10$, $\text{Mg}^{2+} = 0.011$, $\text{K}^+ = 0.004$ mmol l^{-1} , hardness as $\text{CaCO}_3 = 3.3$ mg l^{-1} , $\text{pH} = 7.0$). During this time, fish were fed to satiation daily with commercial pellet food (BioTrout 4.0 mm, Bio-Oregon™, Long-view, WA, USA). Food was withheld for 7 days prior to experimentation. For the in vivo series, fish were anaesthetized and surgery was performed as described below. For the in situ series, fish were sampled either at 7 days of fasting (control), or at 24 h or 48 h after voluntary feeding to satiation. For both series, we measured or calculated P_aO_2 , haemoglobin- O_2 saturation (Hb- O_2 saturation), pH_a , $[\text{HCO}_3^-]_a$, P_aCO_2 , total ammonia (Tamm), and/or P_aNH_3 from blood sampled from the dorsal aorta (DA). Venous drainage was sampled from the subintestinal vein (SIV) and measured for P_vO_2 , Hb- O_2 saturation, pH_v , $[\text{HCO}_3^-]_v$, P_vCO_2 , SIV Tamm, and/or P_vNH_3 . All experiments were approved by the UBC Animal Care Committee (AUP 14–0251 and 18–0271) and conformed to national regulations of the Canada Council for Animal Care.

Cannulation and blood sampling

For both in vivo and in situ series, the same DA and SIV cannulation techniques were performed. Fish were anaesthetized with 0.1–0.2 g l⁻¹ NaOH-neutralized MS-222 (Syndel Laboratories, Parksville, BC). Once anaesthetized, the fish were transferred to an operating table and their gills were continuously irrigated with temperature-controlled anesthetic water (~0.03 g l⁻¹) to achieve stage 5 anaesthesia (McFarland 1959). The DA was cannulated according to the procedure described in Soivio et al. (1972). The SIV was cannulated near the proximal end of the posterior intestine, as described by Cooper et al. (2014). In brief, a 3-cm incision in the midline anterior to the anus was made, the SIV was located, and then the most posterior end of the vessel was tied with 2–0 silk thread to occlude posterior blood flow. About 2 cm of polyethylene tubing-10 (PE10; Clay-Adams™, Becton and Dickinson Co., Franklin Lakes, NJ, USA) was fitted to stretched PE50 tubing and filled with heparinized (150 I.U. ml⁻¹; Sigma-Aldrich, St. Louis, MO, USA) Cortland saline (Wolf 1963). An opening for the cannula in the SIV was made with a hypodermal needle, then the PE10-tipped cannula was pushed forward anteriorly about 2 cm and tightly secured around the vessel using 2–0 surgical silk.

For in vivo series, blood flow through the cannula was checked, which was then flushed with heparinized saline. The cannula was secured to the anal fin and the incision was sutured closed with silk suture. In addition, flared PE240 was inserted down the oesophagus into the stomach and securely sutured through a hole next to the nostril to serve as a feeding tube. Fish were then transferred to black Plexiglas chambers and were left to recover overnight. The chambers were supplied with flow-through 9°C freshwater and were aerated throughout the experiment. After overnight recovery, blood was sampled from both the DA and SIV cannulae while the fish was still in the fasting state, which will be referred to as the “control” from here on. Approximately 0.5 ml of blood was sampled from each cannula by blood pressure-driven and/or gravity driven flow into 0.6-ml heparinized microcentrifuge tubes, taking care to minimize air exposure. Equivalent volumes of saline were immediately injected to replace the blood removed by sampling. Immediately after, fish were fed a 1% body mass ration of pellet food (delivered as a slurry of homogenised food pellets in two volumes of water; Cooper and Wilson 2008; Bucking et al. 2010) via the oesophageal feeding tube. Postprandial blood samples were collected at 4, 8, 12, 24 and 48 h after feeding.

For the in situ series, fish were implanted with the same DA and SIV cannulae on a surgery table with anaesthesia and gill irrigation maintained at the same level throughout. Experiments lasted approximately 1 h. Blood samples were collected via the cannulae into 0.6-ml heparinized

microcentrifuge tubes. In addition, the hepatic portal vein (HPV) blood was sampled by direct insertion of the needle of a heparinized gas-tight 100 µL-syringe (Hamilton, Reno, Nevada, USA) into the blood vessel.

Whole blood measurements

Collected blood samples were first measured for pH, PO₂ and PCO₂ by inserting the micro-electrode and micro-optodes to the bottom of the microcentrifuge tube, while kept at the experimental temperature. The pH was measured using an oesophageal pH microelectrode (MI-508; 1.4–1.6 mm OD) and a flexible micro-reference electrode (MI-402; Microelectrodes Inc., Bedford, NH, USA) connected to a model 220 pH meter (Corning Instruments, Corning, NY, USA). The electrodes were calibrated with precision buffers at the appropriate temperature (Fisher Scientific and Radiometer-Copenhagen, Copenhagen, Denmark).

The PO₂ and PCO₂ were measured using micro-optodes (PreSens Precision Sensing GmbH Regensburg, Germany) positioned into the collection tubes using micro-manipulators (World Precision Instruments, Sarasota, FL, USA). The PO₂ micro-optodes were calibrated with air-equilibrated and sodium sulfite saturated saline kept in gas-tight bottles. The PCO₂ micro-optodes were prototype devices (PreSens 200 001 368) connected to an electronic transmitter (PreSens 300 000 114), with the output displayed on a personal computer running prototype software (PreSens 200 001 488). These PCO₂ micro-optodes were prepared and calibrated as described in Jung et al. (2020), with salines equilibrated to 0.04, 0.5, 1, 3, and 5% CO₂ using CO₂/air mixtures created by a 301aF precision gas-mixing pump (Wösthoff Messtechnik GmbH, Bochum, Germany). All calibration solutions were kept at the experimental temperature of 9 °C.

The haemoglobin-O₂ saturation (%) was calculated from the measured blood O₂ content ([O₂]) and haemoglobin concentration ([Hb]) by taking 4 O₂ molecules per Hb (tetramer) into account. Blood [O₂] was measured using the Tucker method (Tucker, 1967). During collection of the blood from cannulae into the microcentrifuge tubes, 10 µL of blood was taken using a gas-tight syringe and injected into a custom-made 2344 µL Tucker chamber with the PO₂ micro-optode inserted through a gas-tight septum. The chamber was filled with a potassium ferricyanide and saponin solution to release O₂ bound by haemoglobin. The [Hb] was measured as described by Kampen and Zijlstra (1961). After all measurements were completed, the remaining blood was centrifuged (2 min, 5000 g), and the plasma was removed and flash-frozen in liquid N₂ and stored at –70 °C in an ultra-cold freezer.

Plasma and chyme measurements: Tamm and TCO_2

At the end of the experiment, fish were euthanized by an overdose of neutralized MS-222. The four segments of the GIT (stomach, anterior intestine, mid intestine, posterior intestine) were ligated and excised, and the chyme was collected into 2-ml centrifuge tubes. The chyme samples were centrifuged (2 min, 5000 g), and the pH values of the supernatants were measured. The samples were then flash-frozen in liquid N_2 for storage at -70°C in an ultra-cold freezer.

Plasma and chyme supernatant were later measured for total CO_2 content (TCO_2) using a Corning 965 CO_2 analyser (Ciba-Corning Diagnostics, Halstead, Essex, UK) calibrated with NaHCO_3 standards. The samples were also analyzed for total ammonia (Tamm; μM) by an enzymatic assay based on the glutamate dehydrogenase/NAD method using a commercial kit (Raichem; San Diego, CA, USA).

Calculations and statistical analyses

The $[\text{HCO}_3^-]$ values in plasma and chyme, and the PCO_2 values in chyme were derived from measurements of pH and TCO_2 via rearrangements of the Henderson–Hasselbach equation (see Wood et al., 1983) using values for pK' and CO_2 solubility for teleost plasma (Boutilier et al., 1984). Partial pressures of ammonia (PNH_3) of both plasma and chyme were similarly calculated via the Henderson–Hasselbach equation using measured Tamm and pH values with the pK and solubility coefficients reported in Cameron and Heisler (1983).

Graphs were made and statistical analyses were performed using Graphpad Prism software (version 7.0a). Data have been expressed as means \pm SEM (N =number of fish). For both series, comparisons between DA and SIV at each postprandial sampling time were made using two-way ANOVA with Sidak's post hoc test. Comparisons of the measured parameter of either DA or SIV from different postprandial sampling times to that of the control value were done using two-way ANOVA with Dunnett's post hoc test. Tamm and PNH_3 data were log transformed to meet assumptions of homogeneity of variance, and comparison between plasma and chyme Tamm or PNH_3 values were done using one-way ANOVA with Sidak's post hoc test. One-way ANOVA with Tukey's post hoc test were performed to compare chyme pH, $[\text{HCO}_3^-]$, PCO_2 between GIT compartments. Comparison of pH, $[\text{HCO}_3^-]$ or PCO_2 between posterior chyme and SIV were done using two-way ANOVA with Sidak's post hoc test. A significance level of $p < 0.05$ was used in all tests.

Results

Postprandial blood O_2 (in vivo)

Within 4 h of feeding, P_aO_2 in the DA increased significantly from about 100 to 140 mmHg, remained high for about 8 h, then gradually came back down to control level by 24 and 48 h after feeding (Fig. 1A). Feeding had no significant effect on the P_vO_2 in the SIV (25–49 Torr), which




Fig. 1 Postprandial dorsal aorta (DA) and subintestinal vein (SIV) blood A PO_2 (DA parameters indicated with subscript “a” and SIV with subscript “v”; P_aO_2 and P_vO_2 , respectively; Torr; $N=5-9$) and B haemoglobin saturation (%), measured on rainbow trout in vivo. Control represents fasting state prior to feeding which occurred immedi-

ately thereafter. Values are means \pm SEM ($N=4-6$). †Indicates significant difference between DA and SIV values. All SIV PO_2 were significantly lower than DA PO_2 . *Indicates a significant difference from control values for either DA or SIV

remained lower than P_aO_2 throughout the series. The difference between P_aO_2 and P_vO_2 was approximately 64 mmHg in fasting fish (control) then increased to 98 mmHg at 4 h following feeding. Despite these changes and the observed increase in P_aO_2 in response to feeding, the Hb- O_2 saturation did not change in either arterial (DA) or venous (SIV) blood, averaging $77.3 \pm 2.4\%$ and $44 \pm 3.1\%$, respectively (Fig. 1B).

Postprandial blood and chyme pH, $[HCO_3^-]$, PCO_2 (in vivo)

We observed a metabolic alkalosis (also known as an alkaline tide) at 24 h post-feeding where plasma pH_a and $[HCO_3^-]_a$ were elevated significantly relative to the control values (Fig. 2A, B). A 5.6 mM increase in $[HCO_3^-]_a$ at 24 h led to a 0.4 unit increase in pH_a . Both parameters then

came back down to the control level by 48 h. In contrast, the pH_v and $[HCO_3^-]_v$ remained unchanged throughout digestion. DA and SIV values were similar except at 4 and 24 h for pH, and at 24 h for $[HCO_3^-]$. Throughout the experiment, the measured P_vCO_2 values in the SIV were generally higher than P_aCO_2 by approximately 1.5 Torr (Fig. 2C). Both P_aCO_2 and P_vCO_2 increased similarly at 12 h (by ~ 1.7 and 1.4 Torr, respectively), but came back down to control levels by 24 h. Interestingly, in both DA and SIV blood, the PCO_2 remained below 8 Torr throughout digestion.

The cannulated animals were killed for chyme sampling at 48 h post-feeding. Visual inspection showed that most of the chyme was in the posterior intestine at this time, with small amounts in the more anterior sections of the intestine and stomach. Levels of pH and $[HCO_3^-]$ in the chyme collected at 48 h increased from anterior to posterior sections




Fig. 2 Postprandial DA and SIV blood **A** pH (pH_a and pH_v , respectively; $N=5-9$), **B** $[HCO_3^-]$ ($[HCO_3^-]_a$ and $[HCO_3^-]_v$ respectively; mM; $N=4-9$), and **C** PCO_2 (P_aCO_2 and P_vCO_2 respectively; Torr; $N=4-9$), measured on rainbow trout in vivo. See Fig. 1 legend for further details

of the GIT (Table 1). Consequently, the calculated PCO_2 was the highest in the stomach chyme despite the lowest $[\text{HCO}_3^-]$ due to its low pH. The intestinal chyme $[\text{HCO}_3^-]$ was much higher than that of the $[\text{HCO}_3^-]_v$ in the SIV blood plasma (Table 1 and Fig. 2B; $p=0.005$), especially in the posterior intestine where the SIV blood was sampled. However, the pH (8.0 ± 0.1) and PCO_2 values (15.3 ± 7.7 Torr) in the posterior intestinal chyme were not significantly different from those (7.7 ± 0.1 , 5.8 ± 0.4 Torr, respectively) in the SIV blood plasma (Table 1 and Fig. 2A, C; $p=0.14$ and 0.31 , respectively).

Postprandial plasma and chyme ammonia (in vivo and in situ)

In both the in vivo and in situ series, venous (SIV and/or HPV) plasma Tamm values were generally slightly higher than arterial (DA) plasma Tamm (Fig. 3 A&B; denoted by †). Interestingly, both series showed that DA and SIV Tamm did not increase with feeding. In fact, in the in situ

series, SIV Tamm decreased by 52% at 48 h after feeding (Fig. 3B), a trend that was not seen in the in vivo series (Fig. 3A). Using the Henderson-Hasselbalch equation with measured Tamm and pH, we calculated PNH_3 in plasma and chyme in both series (Fig. 4). The DA and SIV plasma PNH_3 were not significantly different from each other, with values of 30–43 μTorr in the in vivo series and 8–13 μTorr in the in situ series. The SIV plasma Tamm and PNH_3 at 48 h were 79–96% lower than posterior chyme Tamm or PNH_3 collected immediately afterwards in both series (Fig. 4). In contrast, HPV Tamm increased by 132% at 24 h after feeding and remained high until 48 h (Fig. 3B).

The chyme Tamm (Fig. 4A, B) was highest in the stomach in both the in vivo and the in situ series (2184 and 2024 μM , respectively), followed by posterior intestine (1027 and 567 μM , respectively), and anterior and mid intestines (495–722 and 114–160 μM , respectively). Reflecting the pH gradient along the GIT (Table 1); however, PNH_3 was the lowest in the stomach in both series, with 0.02–3.1 μTorr . The intestinal chyme PNH_3 gradually increased down the

Table 1 Measured pH, $[\text{HCO}_3^-]$, and calculated PCO_2 of chyme collected in different GIT segments 48 h post-feeding of the rainbow trout in the in vivo series

	Stomach	Anterior intestine	Mid intestine	Posterior intestine
pH	5.3 ± 0.3^x	7.6 ± 0.1^y	7.7 ± 0.1^y	8.0 ± 0.1^y
$[\text{HCO}_3^-]$ (mM)	1.9 ± 0.9^x	19.4 ± 4.9^y	21.0 ± 5.1^y	40.1 ± 5.8^z
PCO_2 (Torr)	125.5 ± 15.8^x	12.0 ± 2.5^y	15.5 ± 7.0^y	15.3 ± 7.7^y

Letters indicate significant difference between different segments for that particular variable. Values are means \pm SEM (N=7–9 for pH; N=4–8 for $[\text{HCO}_3^-]$ and PCO_2)




Fig. 3 Comparison of postprandial plasma Tamm (μM) from DA, SIV, and HPV blood sampled from rainbow trout **A** in vivo (N=4–6) and **B** in situ (N=5–7). Control represents fasting state, feeding

occurred immediately after control sample was taken in (A). See Fig. 1 legend for further details




Fig. 4 Comparison of measured plasma and chyme (**A, B**) Tamm (μM) and calculated (**C, D**) PNH_3 (μTorr) sampled 48 h post feeding in rainbow trout in (**A, C**) the in vivo series ($N=4-7$) and in (**B, D**) the in situ series ($N=5-7$). Plasma Tamm values are reproduced from Fig. 3. In all graphs, left y-axis represents plasma values and right y-axis represents chyme values. Note different scales between left and

right y-axes. Values are means \pm SEM. DA and SIV plasma values were not significantly different from each other in all panels. Upper case letters indicate significant difference between chyme from different GIT compartments. *Indicates a significant difference between SIV plasma and posterior intestine chyme values

tract, with posterior intestine being significantly higher (in vivo 262; in situ 371 μTorr) than all other measurements including plasma values in both series.

Discussion

Here, we investigated potential equilibration of the two respiratory gases, O_2 and CO_2 , and ammonia (total ammonia = $\text{NH}_3 + \text{NH}_4^+$) across the GIT epithelia in rainbow trout by monitoring arterial and venous blood status in vivo following feeding. Overall, feeding had no significant effect on subintestinal venous blood plasma PO_2 , PCO_2 , and ammonia sampled near the posterior intestine, and these were notably different from the posterior intestine chyme values measured in this study and those previously reported. Thus, our

data support our overall hypotheses that there is minimal equilibration between the lumen and the blood circulation, and that blood gas levels in the venous drainage of the intestine remain largely unchanged after feeding. However, we did observe a significant increase in HPV plasma ammonia level after feeding, which agrees with our hypothesis that ammonia absorption occurs in the anterior part of the GIT.

Postprandial blood O_2 (in vivo)

The control P_aO_2 measured in this study is similar to previously reported values of fish starved for 48 h or more (Tetens and Lykkeboe 1985; Perry and Reid 1992). Feeding significantly increased this value within 4 h, which then gradually dropped back to the control level by 48 h (Fig. 1A; open circles). Interestingly, the arterial blood Hb- O_2 saturation (%)

was not affected by feeding (Fig. 1B) and remained similar to previously measured values of fasted animals (Milligan and Wood 1987) and within the typical range of rainbow trout blood (Rummer and Brauner 2015). This suggests a ventilatory response to feeding, rather than an increase in O_2 carrying capacity or blood- O_2 affinity. This is an opposite response to that known in air-breathing terrestrial vertebrates that reduce ventilation to retain CO_2 , thereby triggering a rise in P_aCO_2 following feeding so as to minimize the rise in blood pH (reviewed by Wang et al. 2001). In water-breathing fish, for which O_2 solubility in the media is much less than CO_2 solubility, it has been understood that hypoventilation would detrimentally compromise O_2 uptake at a time when metabolic demand (MO_2) is elevated during digestion. Thus, hypoventilation is assumed to be an impractical response to feeding in fish, but a detailed study is lacking.

Although P_aO_2 increased after feeding, SIV P_vO_2 and Hb- O_2 saturation did not change throughout digestion (Fig. 1; closed circles). This agrees with our initial hypothesis where we expected to see no effect of low luminal PO_2 on P_vO_2 . The lumen of the rainbow trout GIT remains nearly anoxic (< 1 Torr) even during digestion (Jung et al. 2022a). The lumen of the mammalian GIT also remains virtually anoxic (Kurbel et al. 2006; Bettinger 2015; Kalantar-Zadeh et al. 2019), due to the respiration of inhabiting microbes (Espey 2013) and the oxidative chemistry of the GIT fluids (Friedman et al. 2018). In humans, systemic hypoxemia can downregulate many of the GIT transporters (Ward et al. 2014), hindering its functions in assimilation and absorption. The multifunctional GIT of fish (Grosell et al. 2011) also possesses many of the same transporters as in the human GIT, and so may be maintaining its bloodstream PO_2 to avoid such challenges. Thus, mechanisms appear to be present to prevent hypoxia such as increases in blood flow rate. Indeed, to meet the postprandial increase in GIT O_2 requirement, blood flow is increased and redistributed to the GIT (Axelsson et al. 1989, 2000; Axelsson and Fritsche 1991; Thorarensen and Farrell 2006; Eliason et al. 2008). Human colonic cells are shown to survive in an anoxic environment for up to 96 h with a sign of acclimation to the condition such as lowering levels of oxidative stress (Vissenaekens et al. 2019), but similar mechanisms in teleost intestinal cells are as yet unknown. Feeding induces an increase in whole-animal MO_2 by ~96% by 27 h in adult rainbow trout (Eliason et al. 2008), which is likely accompanied by a simultaneous increase in GIT tissue MO_2 (Taylor and Grosell 2009). Although the difference between P_aO_2 and P_vO_2 was greater following feeding (63 Torr in control vs. 88–99 Torr during digestion), there was no change in SIV Hb- O_2 saturation (Fig. 1B). In fish, however, the GIT blood flow can be controlled and quickly increased to supply metabolically active organs at the time of need (Dupont-Prinet et al. 2009; Gräns et al. 2009a, b; Sundh et al. 2018). In various

species including rainbow trout, feeding is shown to induce redistribution of blood flows to the stomach and intestine, with elevations of 72–136% (Axelsson et al. 1989, 2000; Axelsson and Fritsche 1991; Thorarensen and Farrell 2006; Eliason et al. 2008). Thus, the postprandial increase in GIT O_2 requirement could be presumably met by an increase in blood flow regardless of the luminal PO_2 status.

Postprandial blood and chyme pH, $[HCO_3^-]$, and PCO_2 (in vivo)

Under control (fasting) conditions, pH_a , $[HCO_3^-]_a$, and P_aCO_2 were similar or slightly elevated relative to previous findings (e.g. Bucking and Wood 2008; Cooper and Wilson 2008; Milligan and Wood 1987; Perry and Reid 1992; Wood and Eom 2019). This study found a classic postprandial alkaline tide, where arterial pH and $[HCO_3^-]$ increased (Fig. 2A, B) presumably due to acid secretion by the stomach (reviewed by Hersey and Sachs 1995). In both voluntarily and catheter-fed rainbow trout, plasma $[HCO_3^-]$ level increases by ~3–4 mM, causing blood pH to rise by ~0.2–0.3 units (Bucking and Wood 2008; Cooper and Wilson 2008; Bucking et al. 2009). In this study, we found $[HCO_3^-]$ and pH values peaked at 24 h after feeding with elevations of 5.6 mM and 0.4 units, respectively (Fig. 2A, B). The delayed alkaline tide observed in this study (24 h vs 3–12 h; e.g. Bucking et al. 2009; Bucking and Wood 2008; Cooper and Wilson 2008) could be due to the relatively small ration (reviewed by Wood 2019) introduced via the feeding tube (Cooper and Wilson 2008) or delayed digestion due to stress from the surgery (Eliason et al. 2008). For example, a bigger ration size induces greater base excretion, likely to correct for greater blood-acid-base disturbance (Bucking and Wood 2008; Cooper and Wilson 2008; Wood 2019). Moreover, the lower acclimation temperature used here (9 °C) can slow down digestion (Jobling 1981; McCue 2006) relative to the temperatures used in studies mentioned above (10–15.5 °C). Nonetheless, the increase in pH_a and $[HCO_3^-]_a$ were indicative of animals undergoing digestion.

We found no change in P_aCO_2 following feeding (Figs. 1A and 2C), similar to the observations of Bucking and Wood (2008) in voluntarily fed rainbow trout. Our finding and other recent studies suggest fish have a compensatory mechanism for dealing with alkaline tide that is different from the hypoventilatory CO_2 retention strategy of air-breathing terrestrial vertebrates (e.g. Andrade et al. 2004; Busk et al. 2000; Higgins 1914; Wang et al. 2001). Nonetheless, fish are able to minimize blood pH disturbance relative to the postprandial metabolic base load (Wood et al. 2007; Bucking and Wood 2008; Bucking et al. 2010) by excreting HCO_3^- via branchial Cl^-/HCO_3^- exchange mechanisms (Wood et al. 2005, 2007; Tresguerres et al. 2007; Bucking and Wood 2008), as well as via the urine (Bucking

et al. 2010). However, it is interesting that the increase in $[\text{HCO}_3^-]$ seen in DA plasma was not observed in SIV plasma (Fig. 2B). Some of the base load from the stomach is likely excreted on the first pass through the gills, and the remaining fraction is what was seen in our DA samples. This base load was not detected in the SIV sampling site, downstream of the gills and DA sampling site. In seawater fish, the intestine has the capacity to compensate the plasma base load by HCO_3^- secretion into the lumen where it is used for osmoregulation (Wilson et al. 2002; Wilson and Grosell 2003; Grosell and Genz 2006; Grosell et al. 2007; Taylor et al. 2010) and possibly even for postprandial HCO_3^- excretion in rectal fluid and feces (Wilson et al. 1996; Taylor and Grosell 2006; Bucking et al. 2009). Presumably, the intestine in freshwater fish may also be a potential site for postprandial base excretion to alleviate the alkaline tide in addition to export by the gills and kidney. However, an in vitro study found that freshwater rainbow trout intestine tissue had minimal involvement in removing a metabolic base load (Bucking et al. 2009). Additional studies are required to evaluate this possibility further.

Here, we have recorded the first in vivo measurements of PCO_2 in the SIV blood during digestion (Fig. 2C; closed circles). During the period of active digestion, luminal PCO_2 is found to be very high (calculated values of 25–111 Torr and direct measurements of 20–40 Torr; Jung et al. 2022a, b; Wood and Eom 2019). In mammals, microbial metabolic production, reaction of gastric HCl with food carbonates, and endogenous HCO_3^- secretion contribute to high luminal PCO_2 (Steggerda 1968; Altman 1986; Tomlin et al. 1991; Suarez et al. 1997; Kurbel et al. 2006). In fish, endogenous HCO_3^- secretion by the intestinal epithelium for neutralizing chyme (Bucking and Wood 2009; Wood and Eom 2019; Goodrich et al. 2020) and for osmoregulatory purposes, particularly in seawater fish, can contribute to the high luminal PCO_2 . We found that luminal PCO_2 remained elevated even at 48 h after feeding, and at particularly high levels in the stomach in parallel to its low pH (Table 1).

Until now, it has been difficult to conclude whether such high luminal PCO_2 will equilibrate with the venous bloodstream draining the GIT. In seawater-acclimated rainbow trout (presumably) fasted for 72 h, calculated SIV PCO_2 was low at 4.7 Torr (Cooper et al. 2014). Moreover, in Atlantic salmon fed 24–48 h prior to measurement, calculated HPV PCO_2 was low at 4.8 Torr and remained low at 2.9–3.6 Torr for 7 days (Eliason et al. 2007), implying that PCO_2 may not be equilibrating into the blood draining the stomach as well. Both Jung et al. (2022a, b) and Wood and Eom (2019) found highly variable measured SIV PCO_2 levels (11–27 Torr, 2–34 Torr, respectively) using anaesthetized fed fish sampled in situ, even with direct insertion of the PCO_2 micro-optode into the SIV. The present study was able to measure SIV PCO_2 in non-anaesthetized fish and found

low and constant levels during digestion, similar to the control fasting values. There was a slight elevation at 12 h, but in general, all measurements were below 8 mmHg, much lower than previously reported levels measured in the lumen. The calculated PCO_2 of chyme collected at 48 h following feeding was not significantly different from measured SIV plasma PCO_2 , but we suspect the animals to be near completion of digestion at the time of sampling. In fact, our calculated intestinal chyme PCO_2 levels (12–15.5 Torr; Table 1) are similar to those directly measured in fasted rainbow trout (7–13 Torr; Wood and Eom 2019). Thus, despite the high luminal PCO_2 found during active digestion, this study and previous reported values support that minimal equilibration of PCO_2 is occurring across the GIT epithelium.

Studies on higher vertebrates provide some insights into possible reasons for this lack of PCO_2 equilibration. In mammals, some of the luminal CO_2 is absorbed to the blood, breathed out (Ghoos et al. 1993), removed by microbes, and/or released as flatulence (Ohashi et al. 2007; Lacy et al. 2011; Oliphant and Allen-Vercoe 2019). To large extent, the luminal epithelia in higher vertebrates appear to have limited permeability allowing only a small percentage of other diffusible gases such as H_2 , He and CH_4 to diffuse from the lumen into the villus blood (Bond et al. 1977). As a result, feeding resulted in no change in arterial PCO_2 in humans (Mensink et al. 2006) and no difference between arterial and venous PCO_2 in chickens (*Gallus gallus domesticus*; Wideman et al. 2003). Moreover, mammals possess a countercurrent arrangement of the capillary bed in the intestinal villi and in the ruminant stomach that may possibly function to recirculate and trap the CO_2 . While the rainbow trout do not have intestinal villi, they have larger epithelial folds that are more prominent in the posterior intestine (Harder 1975; Kapoor et al. 1975). If a similar system is present in the intestinal folds of trout as in the intestinal villi of mammals, the luminal PCO_2 could diffuse into and recirculate within the intestinal fold, without appearing in the venous drainage from the intestine. This locally high PCO_2 in the intestinal folds possibly could help with oxygen unloading to the enterocytes via the Bohr and Root effects (Nikinmaa 2006; Cooper et al. 2014; Rummer and Brauner 2015). Thus, it will be of great interest to further investigate the vascular micro-anatomy of the trout GIT in the future.

Postprandial plasma ammonia (in vivo and in situ)

In both the in vivo and the in situ series, control fish had DA plasma Tamm concentrations typical of those measured in other studies (Fig. 3A, B; Bucking et al. 2009; Karlsson et al. 2006; Knoph and MåsØval 1996). The Tamm levels in DA plasma sampled from fish in situ was higher than in vivo, which could be due to anaesthetic in the water reducing the ammonia excretion rate (Guo et al. 1995). More significantly,

SIV Tamm did not increase after feeding in both series, supporting our initial hypothesis. Our chyme measurements are in agreement with earlier studies (Bucking and Wood 2012; Rubino et al. 2014; Jung et al. 2022a) showing that chyme Tamm throughout the intestine is typically very high (Fig. 4A, B; 495–1027 μM in vivo; 114–566 μM in situ) following feeding in rainbow trout. This likely reflects the catabolism of amino acids within the GIT lumen after ingestion of protein. Importantly, these data also confirm our recent finding (Jung et al. 2022a) that chyme Tamm levels are even higher in the stomach (2024–2181 μM) than in the more posterior sections of the GIT (Fig. 4A, B). This suggests considerable absorption of ammonia in the anterior tract, perhaps in the stomach itself and/or anterior intestine. In fact, rainbow trout stomach tissue in vitro showed the greatest absorption and consumption of ammonia compared to the intestinal tissues (Jung et al. 2022b). And many studies have shown that plasma ammonia level increases up to threefold after feeding (Kaushik and de Oliva Teles 1985; Wicks and Randall 2002; Bucking and Wood 2008, 2012). These previous studies have measured ammonia levels in the mixed arterial–venous plasma samples collected from caudal punctures. With our localized blood sampling method, we were able to show that any postprandial ammonia absorption that occurred in the GIT in our study was likely metabolized in the liver and/or excreted to the water via the gills (Bucking and Wood 2008), thus resulting in no change in arterial plasma ammonia levels. However, we cannot eliminate the possibility that the small ration size used in our study contributed to this lack of elevation.

Previous studies found that feeding stimulates the endogenous ammonia production rate of the intestinal epithelia and increases the efflux rate of ammonia to the serosal solution in vitro (Rubino et al. 2014; Jung et al. 2021). Whole-body ammonia excretion rate increases after feeding (Alsop and Wood 1997; Taylor et al. 2007; Bucking et al. 2010), of which up to 47% can originate from the GIT, mainly by endogenous production (Rubino et al. 2014). However, we measured 5–sevenfold lower Tamm in venous drainage relative to the posterior intestine chyme (Fig. 4A, B). With measured Tamm and pH, we calculated PNH_3 of both plasma and chyme (Fig. 4C, D). Although our chyme PNH_3 values were lower than reported values at an earlier stage of the digestion (24 h after feeding; Jung et al. 2022a, b; Rubino et al. 2014), they were still significantly higher than SIV PNH_3 in both series. Thus, like PCO_2 (Sect. "Postprandial blood and chyme pH, $[\text{HCO}_3^-]$, and PCO_2 (in vivo)"), there was minimal equilibration of PNH_3 between the lumen and SIV.

In contrast, there was a clear increase in HPV Tamm sampled in situ (Fig. 3B), similar to findings by Karlsson et al. (2006). The HPV differs from the SIV by additionally

collecting venous drainage from the stomach and anterior intestine, suggesting that the anterior part of the GIT could be a greater site of ammonia absorption. In fact, Rubino et al. (2019) have presented evidence that NKCC functions as an NH_4^+ transporter in the anterior intestine of trout, as it does in mammalian gastrointestinal tissues, probably by substitution of NH_4^+ at the K^+ site on the co-transporter (Abdoun et al. 2006; Worrell et al. 2008). Furthermore, Rubino et al. (2019) showed that NKCC is more highly expressed in the anterior intestine than in the mid and posterior intestine of fed rainbow trout. Another possible ammonia NH_3 transporter—rhesus glycoprotein (Rh protein)—is expressed along the mice GIT including the stomach; both apical (Rhcg) and basolateral (Rhbg) isoforms are present throughout the tract (Handlogten et al. 2005). In rainbow trout, the Rhbg is expressed along the whole GIT including the stomach, with feeding elevating the relative expression in both the anterior and posterior intestines (Bucking and Wood 2012). The Rhcg seems to be lacking in the intestine of rainbow trout (Nawata et al. 2007) and the situation is currently unknown in the stomach. However, two Rhcg isoforms, in addition to Rhbg, were expressed throughout the intestine in two marine Batrachoidid teleosts (Bucking et al. 2013). Nonetheless, our data suggest that the posterior intestine is not the major source of postprandial ammonia absorption, and that ammonia may be transported in the anterior intestine and/or even across the stomach (Jung et al. 2022b) into the portal bloodstream to be processed in the liver. NH_3 absorption across the stomach would be problematical, as it would mean moving against a large PNH_3 gradient (Fig. 4C, D; 0.02–3.1 μTorr in stomach vs 13–30 μTorr in SIV). More likely, ammonia is transported by the gastric epithelium in the ionic form NH_4^+ as there is now considerable evidence that the stomach plays an important role in the absorption of other cations (Na^+ , K^+ , Ca^{2+} , Mg^{2+}) in the rainbow trout (Bucking and Wood 2006, 2007; Bucking et al. 2011).

In summary, it is now clear that feeding and digestive processes establish a harsh environment in the GIT lumen of the rainbow trout: high PCO_2 (Wood and Eom 2019; Jung et al. 2020, 2022a), high ammonia (Jung et al. 2022a), and virtual anoxia (Jung et al. 2020, 2022a). This study found that despite such conditions and increased blood flow to the system after feeding (Eliason et al. 2008), the composition of the venous blood draining the posterior intestine remains relatively unchanged, indicating minimal equilibration of PO_2 , PCO_2 and ammonia.

Acknowledgements We thank Dr. Junho Eom for his assistance with cannulation and PreSens GmbH (Regensburg, Germany) for the loan of their PCO_2 measurement system.

Funding This study was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grants (NSERC RGPIN 2017-03843 to CMW and RGPIN-2018-04172 to CJB).

Data availability The datasets generated and analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Conflict of interest We declare we have no competing interests.

References

- Abdoun K, Stumpff F, Martens H (2006) Ammonia and urea transport across the rumen epithelium: a review. *Anim Health Res Rev* 7:43–59. <https://doi.org/10.1017/S1466252307001156>
- Alabaster JS, Herbert DWM, Hemens J (1957) The survival of rainbow trout (*Salmo Gairdnerii* Richardson) and perch (*Perca Fluvialtilis* L.) at various concentrations of dissolved oxygen and carbon dioxide. *Ann Appl Biol* 45:177–188. <https://doi.org/10.1111/j.1744-7348.1957.tb00452.x>
- Alsop DH, Wood CM (1997) The interactive effects of feeding and exercise on oxygen consumption, swimming performance and protein usage in juvenile rainbow trout (*Oncorhynchus mykiss*). *J Exp Biol* 200:2337–2346
- Altman F (1986) Downwind update—a discourse on matters gaseous. *West J Med* 145:502–505
- Andrade DV, De Toledo LF, Abe AS, Wang T (2004) Ventilatory compensation of the alkaline tide during digestion in the snake *Boa constrictor*. *J Exp Biol* 207:1379–1385. <https://doi.org/10.1242/jeb.00896>
- Axelsson M, Fritsche R (1991) Effects of exercise, hypoxia and feeding on the gastrointestinal blood flow in the Atlantic cod *Gadus morhua*. *J Exp Biol* 158:181–198
- Axelsson M, Driedzic WR, Farrell AP, Nilsson S (1989) Regulation of cardiac output and gut blood flow in the sea raven, *Hemirhamphus americanus*. *Fish Physiol Biochem* 6:315–326
- Axelsson M, Thorarensen H, Nilsson S, Farrell AP (2000) Gastrointestinal blood flow in the red Irish lord, *Hemilepidotus hemilepidotus*: long-term effects of feeding and adrenergic control. *J Comp Physiol B* 170:145–152. <https://doi.org/10.1007/s003600050269>
- Beamish FWH, Thomas E (1984) Effects of dietary protein and lipid on nitrogen losses in rainbow trout, *Salmo gairdneri*. *Aquaculture* 41:359–371. [https://doi.org/10.1016/0044-8486\(84\)90203-5](https://doi.org/10.1016/0044-8486(84)90203-5)
- Bettinger CJ (2015) Materials advances for next-generation ingestible electronic medical devices. *Trends Biotechnol* 33:575–585. <https://doi.org/10.1016/j.tibtech.2015.07.008>
- Bond JH, Levitt DG, Levitt MD (1977) Quantitation of countercurrent exchange during passive absorption from the dog small intestine. *J Clin Invest* 59:308–318. <https://doi.org/10.1172/JCI108642>
- Bucking C, Wood CM (2006) Gastrointestinal processing of Na⁺, Cl⁻, and K⁺ during digestion: implications for homeostatic balance in freshwater rainbow trout. *Am J Physiol-Regul, Integr Comp Physiol* 291:R1764–R1772. <https://doi.org/10.1152/ajpregu.00224.2006>
- Bucking C, Wood CM (2007) Gastrointestinal transport of Ca²⁺ and Mg²⁺ during the digestion of a single meal in the freshwater rainbow trout. *J Comp Physiol B* 177:349–360. <https://doi.org/10.1007/s00360-006-0134-3>
- Bucking C, Wood CM (2008) The alkaline tide and ammonia excretion after voluntary feeding in freshwater rainbow trout. *J Exp Biol* 211:2533–2541. <https://doi.org/10.1242/jeb.015610>
- Bucking C, Wood CM (2009) The effect of postprandial changes in pH along the gastrointestinal tract on the distribution of ions between the solid and fluid phases of chyme in rainbow trout. *Aquac Nutr* 15:282–296. <https://doi.org/10.1111/j.1365-2095.2008.00593.x>
- Bucking C, Wood CM (2012) Digestion of a single meal affects gene expression of ion and ammonia transporters and glutamine synthetase activity in the gastrointestinal tract of freshwater rainbow trout. *J Comp Physiol B* 182:341–350. <https://doi.org/10.1007/s00360-011-0622-y>
- Bucking C, Fitzpatrick JL, Nadella SR, Wood CM (2009) Post-prandial metabolic alkalosis in the seawater-acclimated trout: the alkaline tide comes in. *J Exp Biol* 212:2159–2166. <https://doi.org/10.1242/jeb.027862>
- Bucking C, Landman MJ, Wood CM (2010) The role of the kidney in compensating the alkaline tide, electrolyte load, and fluid balance disturbance associated with feeding in the freshwater rainbow trout, *Oncorhynchus mykiss*. *Comp Biochem Physiol-Mol Integr Physiol* 156:74–83. <https://doi.org/10.1016/j.cbpa.2009.12.021>
- Bucking C, Fitzpatrick JL, Nadella SR et al (2011) Assimilation of water and dietary ions by the gastrointestinal tract during digestion in seawater-acclimated rainbow trout. *J Comp Physiol B* 181:615–630. <https://doi.org/10.1007/s00360-011-0550-x>
- Bucking C, Edwards SL, Tickle P et al (2013) Immunohistochemical localization of urea and ammonia transporters in two congeneric fish species, the ureotelic gulf toadfish (*Opsanus beta*) and the ammoniotelic plainfin midshipman (*Porichthys notatus*). *Cell Tissue Res* 352:623–637. <https://doi.org/10.1007/s00441-013-1591-0>
- Busk M, Overgaard J, Hicks JW et al (2000) Effects of feeding on arterial blood gases in the American alligator *Alligator mississippiensis*. *J Exp Biol* 203:3117–3124
- Cameron JN, Heisler N (1983) Studies of ammonia in the rainbow trout: physico-chemical parameters, acid-base behaviour and respiratory clearance. *J Exp Biol* 105:107–125
- Cooper CA, Wilson RW (2008) Post-prandial alkaline tide in freshwater rainbow trout: effects of meal anticipation on recovery from acid-base and ion regulatory disturbances. *J Exp Biol* 211:2542–2550. <https://doi.org/10.1242/jeb.015586>
- Cooper CA, Regan MD, Brauner CJ et al (2014) Osmoregulatory bicarbonate secretion exploits H⁺-sensitive haemoglobins to autoregulate intestinal O₂ delivery in euryhaline teleosts. *J Comp Physiol B* 184:865–876. <https://doi.org/10.1007/s00360-014-0844-x>
- de Solbé JFLG, Shurben DG (1989) Toxicity of ammonia to early life stages of rainbow trout (*Salmo gairdneri*). *Water Res* 23:127–129. [https://doi.org/10.1016/0043-1354\(89\)90071-7](https://doi.org/10.1016/0043-1354(89)90071-7)
- Dupont-Prinet A, Claireaux G, McKenzie DJ (2009) Effects of feeding and hypoxia on cardiac performance and gastrointestinal blood flow during critical speed swimming in the sea bass *Dicentrarchus labrax*. *Comp Biochem Physiol-Mol Integr Physiol* 154:233–240. <https://doi.org/10.1016/j.cbpa.2009.06.015>
- Eliason EJ, Kiessling A, Karlsson A et al (2007) Validation of the hepatic portal vein cannulation technique using Atlantic salmon *Salmo salar* L. *J Fish Biol* 71:290–297. <https://doi.org/10.1111/j.1095-8649.2007.01489.x>
- Eliason EJ, Higgs DA, Farrell AP (2008) Postprandial gastrointestinal blood flow, oxygen consumption and heart rate in rainbow trout (*Oncorhynchus mykiss*). *Comp Biochem Physiol-Mol Integr Physiol* 149:380–388. <https://doi.org/10.1016/j.cbpa.2008.01.033>
- Ellis RP, Urbina MA, Wilson RW (2017) Lessons from two high CO₂ worlds – future oceans and intensive aquaculture. *Glob Chang Biol* 23:2141–2148. <https://doi.org/10.1111/gcb.13515>
- Espes MG (2013) Role of oxygen gradients in shaping redox relationships between the human intestine and its microbiota. *Free Radic Biol Med* 55:130–140. <https://doi.org/10.1016/j.freeradbiomed.2012.10.554>
- Food and Agriculture Organization (2020) The State of world fisheries and aquaculture. FAO, Rome, Italy
- Friedman ES, Bettinger K, Espipova TV et al (2018) Microbes vs. chemistry in the origin of the anaerobic gut lumen. *Proc Natl Acad Sci U S A* 115:4170–4175. <https://doi.org/10.1073/pnas.1718635115>

- Ghoos YF, Maes BD, Geypens BJ et al (1993) Measurement of gastric emptying rate of solids by means of a carbon-labeled octanoic acid breath test. *Gastroenterology* 104:1640–1647. [https://doi.org/10.1016/0016-5085\(93\)90640-X](https://doi.org/10.1016/0016-5085(93)90640-X)
- Goodrich HR, Bayley M, Birgersson L et al (2020) Understanding the gastrointestinal physiology and responses to feeding in air-breathing Anabantiform fishes. *J Fish Biol* 96:986–1003. <https://doi.org/10.1111/jfb.14288>
- Gräns A, Albertsson F, Axelsson M, Olsson C (2009a) Postprandial changes in enteric electrical activity and gut blood flow in rainbow trout (*Oncorhynchus mykiss*) acclimated to different temperatures. *J Exp Biol* 212:2550–2557. <https://doi.org/10.1242/jeb.030593>
- Gräns A, Axelsson M, Pitsillides K et al (2009b) A fully implantable multi-channel biotelemetry system for measurement of blood flow and temperature: a first evaluation in the green sturgeon. *Hydrobiologia* 619:11–25. <https://doi.org/10.1007/s10750-008-9578-7>
- Grosell M, Genz J (2006) Ouabain-sensitive bicarbonate secretion and acid absorption by the marine teleost fish intestine play a role in osmoregulation. *Am J Physiol Regul Integr Comp Physiol* 291:R1145–R1156. <https://doi.org/10.1152/ajpregu.00818.2005>
- Grosell M, Gilmour KM, Perry SF (2007) Intestinal carbonic anhydrase, bicarbonate, and proton carriers play a role in the acclimation of rainbow trout to seawater. *Am J Physiol-Regul, Integr Comp Physiol* 293:R2099–R2111. <https://doi.org/10.1152/ajpregu.00156.2007>
- Grossell M, Farrell AP, Brauner CJ (2011) *The multifunctional gut of fish*. Elsevier Inc
- Guo F, Teo L, Chen T (1995) Effects of anaesthetics on the water parameters in a simulated transport experiment of platyfish, *Xiphophorus maculatus* (Günther). *Aquac Res* 26:265–271. <https://doi.org/10.1111/j.1365-2109.1995.tb00911.x>
- Handlogten ME, Hong SP, Zhang L et al (2005) Expression of the ammonia transporter proteins Rh B glycoprotein and Rh C glycoprotein in the intestinal tract. *Am J Physiol Gastrointest Liver Physiol* 288:1036–1047. <https://doi.org/10.1152/ajpgi.00418.2004>
- Harder W (1975) *Anatomy of fishes*. Schweizerbart, E. Schweizerbart'sche Verlagsbuchhandlung
- Hersey SJ, Sachs G (1995) Gastric acid secretion. *Physiol Rev* 75:155–189. <https://doi.org/10.1152/physrev.1995.75.1.155>
- Higgins HL (1914) The influence of food, posture, and other factors on the alveolar carbon dioxide tension in man. *Am J Physiol-Leg Content* 34:114–126. <https://doi.org/10.1152/ajplegacy.1914.34.1.114>
- Jobling M (1981) The influences of feeding on the metabolic rate of fishes: a short review. *J Fish Biol* 18:385–400. <https://doi.org/10.1111/j.1095-8649.1981.tb03802.x>
- Jung EH, Eom J, Brauner CJ et al (2020) The gaseous gastrointestinal tract of a seawater teleost, the English sole (*Parophrys vetulus*). *Comp Biochem Physiol-Mol Integr Physiol* 247:110743. <https://doi.org/10.1016/j.cbpa.2020.110743>
- Jung EH, Smich J, Rubino JG, Wood CM (2021) An *in vitro* study of urea and ammonia production and transport by the intestinal tract of fed and fasted rainbow trout: responses to luminal glutamine and ammonia loading. *J Comp Physiol B*. <https://doi.org/10.1007/s00360-020-01335-9>
- Jung EH, Brauner CJ, Wood CM (2022a) Post-prandial respiratory gas and acid-base profiles in the gastrointestinal tract and its venous drainage in freshwater rainbow trout (*Oncorhynchus mykiss*) and seawater English sole (*Parophrys vetulus*). *Comp Biochem Physiol-Mol Integr Physiol* 265:111123. <https://doi.org/10.1016/j.cbpa.2021.111123>
- Jung EH, Nguyen J, Nelson C et al (2022b) Ammonia transport is independent of PNH₃ gradients across the gastrointestinal epithelia of the rainbow trout: a role for the stomach. *J Exp Zool A Ecol Integr Physiol*. <https://doi.org/10.1002/jez.2670>
- Kalantar-Zadeh K, Berean KJ, Burgell RE et al (2019) Intestinal gases: influence on gut disorders and the role of dietary manipulations. *Nat Rev Gastroenterol Hepatol* 16:733–747. <https://doi.org/10.1038/s41575-019-0193-z>
- Kampen EJ, Zijlstra WG (1961) Standardization of hemoglobinometry II. The hemiglobincyanide method. *Clin Chim Acta* 6:538–544. [https://doi.org/10.1016/0009-8981\(61\)90145-0](https://doi.org/10.1016/0009-8981(61)90145-0)
- Kapoor BG, Smit HJ, Verighina IA (1975) The alimentary canal and digestion in teleosts. *Adv Mar Biol* 13:109–239
- Karlsson A, Eliason EJ, Mydland LT et al (2006) Postprandial changes in plasma free amino acid levels obtained simultaneously from the hepatic portal vein and the dorsal aorta in rainbow trout (*Oncorhynchus mykiss*). *J Exp Biol* 209:4885–4894. <https://doi.org/10.1242/jeb.02597>
- Kaushik SJ, de Oliva TA (1985) Effect of digestible energy on nitrogen and energy balance in rainbow trout. *Aquaculture* 50:89–101. [https://doi.org/10.1016/0044-8486\(85\)90155-3](https://doi.org/10.1016/0044-8486(85)90155-3)
- Knoph MB, Måsøval K (1996) Plasma ammonia and urea levels in Atlantic salmon farmed in sea water. *J Fish Biol* 49:165–168. <https://doi.org/10.1006/jfbi.1996.0144>
- Kurbel S, Kurbel B, Včev A (2006) Intestinal gases and flatulence: possible causes of occurrence. *Med Hypotheses* 67:235–239. <https://doi.org/10.1016/j.mehy.2006.01.057>
- Lacy BE, Gabbard SL, Crowell MD (2011) Pathophysiology, evaluation, and treatment of bloating: Hope, hype, or hot air? *Gastroenterol Hepatol (n Y)* 7:729–739
- McCue MD (2006) Specific dynamic action: a century of investigation. *Comp Biochem Physiol-Mol Integr Physiol* 144:381–394. <https://doi.org/10.1016/j.cbpa.2006.03.011>
- McFarland WN (1959) A study of the effects of anaesthetics on the behaviour and physiology of fishes. *Publ Inst Mar Sci* 6:22–55
- Mensink PBF, Geelkerken RH, Huisman AB et al (2006) Effect of various test meals on gastric and jejunal carbon dioxide: a study in healthy subjects. *Scand J Gastroenterol* 41:1290–1298. <https://doi.org/10.1080/0036520600670059>
- Milligan CL, Wood CM (1987) Regulation of blood oxygen transport and red cell pHi after exhaustive activity in rainbow trout (*Salmo gairdneri*) and starry flounder (*Platichthys stellatus*). *J Exp Biol* 133:263–282
- Nawata CM, Hung CCY, Tsui TKN et al (2007) Ammonia excretion in rainbow trout (*Oncorhynchus mykiss*): evidence for Rh glycoprotein and H⁺-ATPase involvement. *Physiol Genomics* 31:463–474. <https://doi.org/10.1152/physiolgenomics.00061.2007>
- Nikinmaa M (2006) Gas transport. In: Evans DH, Claiborne JB (eds) *The physiology of fishes*, 3rd edn. CRC Press, Boca Raton, pp 153–174
- Ohashi Y, Igarashi T, Kumazawa F, Fujisawa T (2007) Analysis of acetogenic bacteria in human feces with formyltetrahydrofolate synthetase sequences. *Biosci Microflora* 26:37–40. <https://doi.org/10.12938/bifidus.26.37>
- Oliphant K, Allen-Vercoe E (2019) Macronutrient metabolism by the human gut microbiome: major fermentation by-products and their impact on host health. *Microbiome* 7:1–15. <https://doi.org/10.1186/s40168-019-0704-8>
- Perry SF, Gilmour KM (2006) Acid-base balance and CO₂ excretion in fish: unanswered questions and emerging models. *Respir Physiol Neurobiol* 154:199–215. <https://doi.org/10.1016/j.resp.2006.04.010>
- Perry SF, Reid SD (1992) Relationship between blood O₂ content and catecholamine levels during hypoxia in rainbow trout and American eel. *Am J Physiol* 263:R240–R249
- Randall DJ, Tsui TKN (2002) Ammonia toxicity in fish. *Mar Pollut Bull* 45:17–23
- Rubino JG, Zimmer AM, Wood CM (2014) An *in vitro* analysis of intestinal ammonia handling in fasted and fed freshwater rainbow

- trout (*Oncorhynchus mykiss*). J Comp Physiol B 184:91–105. <https://doi.org/10.1007/s00360-013-0781-0>
- Rubino JG, Wilson JM, Wood CM (2019) An *in vitro* analysis of intestinal ammonia transport in fasted and fed freshwater rainbow trout: roles of NKCC, K⁺ channels, and Na⁺, K⁺ ATPase. J Comp Physiol B 189:549–566. <https://doi.org/10.1007/s00360-019-01231-x>
- Rummer JL, Brauner CJ (2015) Root effect haemoglobins in fish may greatly enhance general oxygen delivery relative to other vertebrates. PLoS ONE 10:1–17. <https://doi.org/10.1371/journal.pone.0139477>
- Soivio A, Westman K, Nyholm K (1972) Improved method of dorsal aorta catheterization: hematological effects followed for three weeks in rainbow trout (*Salmo gairdneri*). Finn Fish Res 1:11–21
- Steggerda FR (1968) Gastrointestinal gas following food consumption. Ann N Y Acad Sci 150:57–66. <https://doi.org/10.1111/j.1749-6632.1968.tb19031.x>
- Suarez F, Furne J, Springfield J, Levitt M (1997) Insights into human colonic physiology obtained from the study of flatus composition. Am J Physiol 272:G1028–G1033. <https://doi.org/10.1152/ajpgi.1997.272.5.G1028>
- Sundh H, Gräns A, Brijs J et al (2018) Effects of coeliacomesenteric blood flow reduction on intestinal barrier function in rainbow trout *Oncorhynchus mykiss*. J Fish Biol 93:519–527. <https://doi.org/10.1111/jfb.13658>
- Taylor JR, Grosell M (2006) Feeding and osmoregulation: dual function of the marine teleost intestine. J Exp Biol 209:2939–2951. <https://doi.org/10.1242/jeb.02342>
- Taylor JR, Grosell M (2009) The intestinal response to feeding in seawater gulf toadfish, *Opsanus beta*, includes elevated base secretion and increased epithelial oxygen consumption. J Exp Biol 212:3873–3881. <https://doi.org/10.1242/jeb.034579>
- Taylor JR, Whittamore JM, Wilson RW, Grosell M (2007) Postprandial acid-base balance and ion regulation in freshwater and seawater-acclimated European flounder, *Platichthys flesus*. J Comp Physiol B 177:597–608. <https://doi.org/10.1007/s00360-007-0158-3>
- Taylor JR, Mager EM, Grosell M (2010) Basolateral NBCe1 plays a rate-limiting role in transepithelial intestinal HCO₃⁻ secretion, contributing to marine fish osmoregulation. J Exp Biol 213:459–468. <https://doi.org/10.1242/jeb.029363>
- Tetens V, Lykkeboe G (1985) Acute exposure of rainbow trout to mild and deep hypoxia: O₂ affinity and O₂ capacitance of arterial blood. Respir Physiol 61:221–235. [https://doi.org/10.1016/0034-5687\(85\)90128-8](https://doi.org/10.1016/0034-5687(85)90128-8)
- Thomas S, Fritsche R, Perry SF (1994) Pre- and post-branchial blood respiratory status during acute hypercapnia or hypoxia in rainbow trout, *Oncorhynchus mykiss*. J Comp Physiol B 164:451–458. <https://doi.org/10.1007/BF00714582>
- Thorarensen H, Farrell AP (2006) Postprandial intestinal blood flow, metabolic rates, and exercise in Chinook salmon (*Oncorhynchus tshawytscha*). Physiol Biochem Zool 79:688–694. <https://doi.org/10.1086/505512>
- Tomlin J, Lewis C, Read NW, Tomlin SJ (1991) Investigation of normal flatus production in healthy volunteers. Gut 32:665–669
- Tresguerres M, Parks SK, Wood CM, Goss GG (2007) V-H⁺-ATPase translocation during blood alkalosis in dogfish gills: interaction with carbonic anhydrase and involvement in the postfeeding alkaline tide. Am J Physiol-Regul, Integr Comp Physiol 292:R2012–R2019. <https://doi.org/10.1152/ajpregu.00814.2006>
- Vissenaekens H, Grootaert C, Rajkovic A et al (2019) The response of five intestinal cell lines to anoxic conditions *in vitro*. Biol Cell 111:232–244. <https://doi.org/10.1111/boc.201800076>
- Wang T, Busk M, Overgaard J (2001) The respiratory consequences of feeding in amphibians and reptiles. Comp Biochem Physiol-Mol Integr Physiol 128:533–547. [https://doi.org/10.1016/s1095-6433\(00\)00334-2](https://doi.org/10.1016/s1095-6433(00)00334-2)
- Ward JBJ, Keely SJ, Keely SJ (2014) Oxygen in the regulation of intestinal epithelial transport. J Physiol 592:2473–2489. <https://doi.org/10.1113/jphysiol.2013.270249>
- Wicks BJ, Randall DJ (2002) The effect of sub-lethal ammonia exposure on fed and unfed rainbow trout: the role of glutamine in regulation of ammonia. Comp Biochem Physiol-Mol Integr Physiol 132:275–285. [https://doi.org/10.1016/S1095-6433\(02\)00034-X](https://doi.org/10.1016/S1095-6433(02)00034-X)
- Wideman RF, Hooze DM, Cummings KR (2003) Dietary sodium bicarbonate, cool temperatures, and feed withdrawal: impact on arterial and venous blood-gas values in broilers. Poult Sci 82:560–570. <https://doi.org/10.1093/ps/82.4.560>
- Wilson RW, Grosell M (2003) Intestinal bicarbonate secretion in marine teleost fish - source of bicarbonate, pH sensitivity, and consequences for whole animal acid-base and calcium homeostasis. Biochim Biophys Acta Biomembr. <https://doi.org/10.1016/j.bbamem.2003.09.014>
- Wilson RW, Gilmour KM, Henry RP, Wood CM (1996) Intestinal base excretion in the seawater-adapted rainbow trout: a role in acid-base balance? J Exp Biol 199:2331–2343
- Wilson RW, Wilson JM, Grosell M (2002) Intestinal bicarbonate secretion by marine teleost fish - why and how? Biochim Et Biophys Acta-Biomembr 1566:182–193. <https://doi.org/10.1016/j.bbamem.2005.01.004>
- Wolf K (1963) Physiological salines for fresh water teleosts. Progress Fish Cult 25:135–140
- Wood CM (2019) Internal spatial and temporal CO₂ dynamics: fasting, feeding, drinking, and the alkaline tide. In: Grosell M, Munday PL, Farrell AP, Brauner CJ (eds) Fish physiology: carbon dioxide. Elsevier Inc., pp 245–286
- Wood CM, Eom J (2019) The internal CO₂ threat to fish: high PCO₂ in the digestive tract. Proc R Soc B Biol Sci. <https://doi.org/10.1098/rspb.2019.0832>
- Wood CM, Kajimura M, Mommsen TP, Walsh PJ (2005) Alkaline tide and nitrogen conservation after feeding in an elasmobranch (*Squalus acanthias*). J Exp Biol 208:2693–2705. <https://doi.org/10.1242/jeb.01678>
- Wood CM, Bucking C, Fitzpatrick J, Nadella S (2007) The alkaline tide goes out and the nitrogen stays in after feeding in the dogfish shark, *Squalus acanthias*. Respir Physiol Neurobiol 159:163–170. <https://doi.org/10.1016/j.resp.2007.06.008>
- Worrell RT, Merk L, Matthews JB (2008) Ammonium transport in the colonic crypt cell line, T84: role for rhesus glycoproteins and NKCC1. Am J Physiol Gastrointest Liver Physiol 294:429–440. <https://doi.org/10.1152/ajpgi.00251.2006>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.